



GROUPE D'ÉTUDES ET DE RECHERCHE
EN ANALYSE DES DÉCISIONS

Les Cahiers du GERAD

CITATION ORIGINALE / ORIGINAL CITATION

GERAD HEC Montréal
3000, ch. de la Côte-Sainte-Catherine
Montréal (Québec) Canada H3T 2A7

Tél. : 514 340-6053
Télec. : 514 340-5665
info@gerad.ca
www.gerad.ca

**Counting the Number of Non-Equivalent
Vertex Colorings of a Graph**

A. Hertz
H. Mélot

G-2013-82

November 2013

Counting the Number of Non-Equivalent Vertex Colorings of a Graph

Alain Hertz

*GERAD & Polytechnique Montréal
Montréal (Québec) Canada, H3A 3A7*

`alain.hertz@gerad.ca`

Hadrien Mélot

*Algorithms Lab
Université de Mons
B-7000 Mons, Belgium*

`hadrien.melot@umons.ac.be`

November 2013

Les Cahiers du GERAD

G-2013-82

Copyright © 2013 GERAD

Abstract: We study the number $\mathcal{P}(G)$ of non-equivalent ways of coloring a given graph G . We show some similarities and differences between this graph invariant and the well known chromatic polynomial. Relations with Stirling numbers of the second kind and with Bell numbers are also given. We then determine the value of this invariant for some classes of graphs. We finally study upper and lower bounds on $\mathcal{P}(G)$ for graphs with fixed maximum degree.

Key Words: Non-equivalent colorings, number of colorings, chromatic polynomial.

Résumé: Nous étudions le nombre $\mathcal{P}(G)$ de colorations non-équivalentes des sommets d'un graphe G . Nous montrons quelques similarités et différences entre ce nouvel invariant et le très célèbre polynôme chromatique. Nous indiquons également quelques relations intéressantes entre cet invariant et les nombres de Stirling du deuxième type et les nombres de Bell. Nous déterminons ensuite la valeur de cet invariant pour quelques classes de graphes. Finalement, nous présentons des bornes inférieures et supérieures sur $\mathcal{P}(G)$ pour des graphes ayant un degré maximum fixé.

Acknowledgments: This work was initiated during a visit – partly funded by the F.R.S.-FNRS – of Alain Hertz at the Algorithms Lab (Université de Mons).

1 Introduction

A question which probably sounds familiar for many researchers in graph theory is: what is the number of ways of coloring a given graph G ? For the path P_3 on three vertices, an answer that makes sense is *two* as depicted in Figure 1. Indeed, at least two colors are needed, and there is only one coloring with two colors (the two extremities share the same color while the central vertex has its own color), and only one coloring with three colors (each vertex has its own color).



Figure 1: The 2 non-equivalent colorings of P_3 (using any number of colors).

However, since more than 100 years, the common answer to the above question for P_3 is not *two* but *twelve*. To understand why, we recall the notion of *chromatic polynomial* which was introduced by Birkhoff in an attempt to prove the four-color theorem. In a paper published in 1912 [1] Birkhoff¹ proves that

“The number of ways of coloring a given map M in k colors ($k = 1, 2, \dots$) is given by a polynomial $P(k)$ of degree n , where n is the number of regions in the map M .”

Birkhoff started the study of this polynomial by defining a quantity m_i as “the number of ways of coloring the map by using exactly i colors when mere permutations of the colors are disregarded”. Then, he used this quantity to define

$$m_i \frac{k!}{(k-i)!}$$

as the “number of ways of coloring the given map in exactly i of the k colors, counting two colorings as distinct when they are obtained by a permutation from the other”. Denoting G the planar graph corresponding to the map M , we can therefore define

$$\Pi(G, k) = \sum_{i=1}^k m_i \frac{k!}{(k-i)!}$$

as the number of ways of coloring G with at most k colors, counting two colorings as distinct when they are obtained by a permutation from the other. The same definition also applies for non-planar graphs G . The *chromatic polynomial* is the polynomial of degree n passing by points $(k, \Pi(G, k))$ for $k = 0, 1, \dots, n$. For example, for the path P_3 we have

$$\Pi(P_3, k) = k(k-1)^2.$$

Indeed, $\Pi(P_3, 0) = \Pi(P_3, 1) = 0$; $\Pi(P_3, 2) = 2$ (take for instance the first two colorings in the left column of Figure 2) and $\Pi(P_3, 3) = 12$ as shown in Figure 2.

The number of vertex colorings of a graph G is nowadays commonly interpreted as $\Pi(G, n)$, where n is the number of vertices in G . However, we argue that the quantity m_i defined by Birkhoff, i.e., the number of *non-equivalent* colorings with an *exact* number i of used colors is also of interest. This is especially the case when a set of elements has to be partitioned into a given number of non-empty subsets, subject to some constraints.

In the next section we fix some notations and give a formal definition of the number $\mathcal{P}(G)$ of non-equivalent vertex colorings of a graph G . Similarities and differences between this invariant and the chromatic polynomial are studied in Section 3. In Section 4, we address the problem of computing $\mathcal{P}(G)$, and we give exact values for some particular graphs in Section 5. Then, in Section 6, we prove some bounds on $\mathcal{P}(G)$ for graphs of bounded maximum degree and let other bounds as open problems.

¹We replaced λ used in [1] by k to unify notations, even in the quotes from this paper.




Figure 2: The 12 colorings of P_3 (using 3 colors) as defined by the chromatic polynomial.

2 Notations

For basic notions of graph theory that are not defined here, we refer to Diestel [3]. Let $G = (V, E)$ be a simple undirected graph. We denote by $n = |V|$ the order of G and by $m = |E|$ its size. We write $G \simeq H$ if G and H are two isomorphic graphs.

Let K_n (resp. C_n and P_n) be the *complete graph* (resp. the *cycle* and the *path*) of order n . The *wheel* W_n is the graph of order n obtained by connecting a vertex to all vertices of C_{n-1} . Also, we write $K_{a,b}$ for the complete bipartite graph where a and b are the cardinalities of the two sets of vertices of the bipartition. Finally, let S_n be the *star* on n vertices, that is $K_{1,n-1}$.

Let $N(v)$ denote the neighbors of a vertex v in G . Vertex v is said to be *simplicial* if $N(v)$ induces a clique in G . A graph is *chordal* (or *triangulated*) if every cycle of length larger than 3 has a chord. The *degree* of a vertex v is denoted $d(v)$ (i.e., $d(v) = |N(v)|$). A vertex v is *isolated* if $d(v) = 0$ and is *dominating* if $d(v) = n - 1$. The *maximum degree* of G is denoted $\Delta(G)$.

Let u and v be two vertices in a graph G of order n , we denote $G \setminus uv$ the graph (of order $n - 1$) obtained by identifying (merging) the vertices u and v and, if $uv \in E(G)$, by removing edge uv . Also, if $uv \in E(G)$, we note $G - uv$ the graph obtained from G by removing edge uv , while if $uv \notin E(G)$, the graph $G + uv$ is the graph obtained by adding uv in G . For a vertex v of G , we denote $G - v$ the graph obtained from G by removing v and all its incident edges.

A *vertex coloring* (or simply a coloring in the sequel) is an assignment of colors to the vertices of G . A *proper coloring* is a coloring such that adjacent vertices have different colors. The *chromatic number* $\chi(G)$ of a graph G is the minimum numbers of colors in a proper coloring of G . Two colorings are *equivalent* if they induce the same partition of the vertex set. We define $P(G, k)$ as the number of proper non-equivalent colorings of a graph G that use *exactly* k colors. The total *number* $\mathcal{P}(G)$ of *non-equivalent colorings* of a graph G is then defined as:

$$\mathcal{P}(G) = \sum_{k=\chi(G)}^n P(G, k). \quad (1)$$

As mentioned in the previous section, $\Pi(G, k)$ is the number of proper colorings of a graph G that use at most k colors, counting two non-identical colorings as distinct when they are obtained by a permutation from the other.

3 Similarities and Differences

According to their definitions, $\Pi(G, k)$ and $P(G, k)$ are linked with the following relations:

$$\Pi(G, k) = \sum_{j=\chi(G)}^k \frac{k!}{(k-j)!} P(G, j), \tag{2}$$

and

$$P(G, k) = \frac{\Pi(G, k) - k\Pi(G, k-1)}{k!}. \tag{3}$$

Observe that unlike $P(G, k)$, $\Pi(G, k)$ also counts colorings with strictly less than k colors. Moreover, while $\mathcal{P}(G)$ and $\Pi(G, n)$ might appear as similar concepts (since they both count colorings with at most n colors), they differ in various ways. We have already mentioned that only non-equivalent colorings are counted in $\mathcal{P}(G)$, which means that $\mathcal{P}(G)$ corresponds to the number of partitions of the vertex set of G , taking into account constraints that prevent some pairs of vertices of belonging to the same subset of the partition. To accentuate these differences, observe that if $\Pi(G, n) < \Pi(H, n)$ for two graphs G and H of order n , this does not necessarily imply that $\mathcal{P}(G) < \mathcal{P}(H)$ (and conversely) as shown in Figure 3.




Figure 3: Two graphs G and H with 6 vertices such that $\Pi(G, 6) < \Pi(H, 6)$ and $\mathcal{P}(G) > \mathcal{P}(H)$.

Also, there exist pairs of graphs (G, H) such that $\mathcal{P}(G) = \mathcal{P}(H)$ but $\Pi(G, n) \neq \Pi(H, n)$, and conversely (see examples in Figure 4).




Figure 4: Two pairs of graphs with 5 vertices showing that equality for one way to counts the colorings does not imply equality for the other.

There are also differences at a computational level. Giving a graph G with a dominating vertex v , the following property states that the computation of $\mathcal{P}(G)$ can be reduced to that of $\mathcal{P}(G - v)$. A similar trivial reduction does not hold for $\Pi(G)$.

Property 1 *If a graph G has a dominating vertex v , then $\mathcal{P}(G) = \mathcal{P}(G - v)$.*

Proof. Since v is a dominating vertex, it must have its own color in all colorings of G , which means that the number of proper non-equivalent colorings of G remains the same when v is removed. \square

Conversely, if G is the disjoint union of two graphs G_1 and G_2 , it is easy to compute $\Pi(G_1 \cup G_2)$ by taking product of $\Pi(G_1)$ and $\Pi(G_2)$. However, the following property shows that the computation of $\mathcal{P}(G_1 \cup G_2)$ is more intricate.

Property 2 Let $G = G_1 \cup G_2$ be a graph that is the disjoint union of two graphs G_1 and G_2 . Then,

$$\mathcal{P}(G) = \sum_{k=1}^n \sum_{i=1}^k \sum_{j=0}^i P(G_1, i) P(G_2, k-j) \binom{i}{i-j} \binom{k-j}{i-j} (i-j)!$$

Proof. The first sum on k comes simply from the definition (1) of $\mathcal{P}(G)$. The two inner sums compute $P(G, k)$ as follows. Let $i \leq k$ be the number of colors used for G_1 . Let j be an integer such that $i - j$ represents the number of colors that are used both in G_1 and in G_2 . The value of j can vary from 0 (that is i colors are shared) to i (that is no color are shared). Observe that in order to use exactly k colors for G , G_2 must be colored with exactly $k - j$ colors. Finally, the term $\binom{i}{i-j}$ counts the numbers of ways to choose the $i - j$ shared colors into G_1 , the term $\binom{k-j}{i-j}$ does the same for G_2 and $(i - j)!$ counts all the possible permutations for this shared colors. \square

As a corollary, we get the following result which will be useful in later sections.

Corollary 3 Let $G = K_p \cup K_q$ be the disjoint union of two cliques of sizes p and q such that $p \leq q$. Then,

$$\mathcal{P}(G) = \sum_{k=q}^{p+q} \binom{p}{k-q} \binom{q}{p+q-k} (p+q-k)!$$

Proof. We apply Property 2 knowing that $\mathcal{P}(K_p, i) = 1$ if and only if $i = p$ and $\mathcal{P}(K_q, k - j) = 1$ if and only if $j = k - q$. For all other values of i and j , the inner products being equal to zero. Also, observe that if $k < q$, there are not enough colors for a proper coloring of G . \square

4 Counting the colorings recursively

As for several other algorithms in graph coloring, the *deletion-contraction* rule is a well known method to compute the chromatic polynomial. More precisely, we have:

$$\Pi(G, k) = \Pi(G - uv, k) - \Pi(G \setminus uv, k),$$

where uv is any edge of G , and

$$\Pi(G, k) = \Pi(G + uv, k) + \Pi(G \setminus uv, k),$$

for any pair of distinct vertices u and v such that $uv \notin E(G)$.

These recurrences, which are often called the *Fundamental Reduction Theorem* [4], are also valid to compute $P(G, k)$ and $\mathcal{P}(G)$. More precisely, let u and v be any pair of distinct vertices of G , we have,

$$P(G, k) = P(G - uv, k) - P(G \setminus uv, k), \tag{4}$$

if $uv \in E(G)$, and

$$P(G, k) = P(G + uv, k) + P(G \setminus uv, k), \tag{5}$$

if $uv \notin E(G)$. Similarly, if $uv \in E(G)$, we have,

$$\mathcal{P}(G) = P(G - uv) - P(G \setminus uv), \tag{6}$$

and

$$\mathcal{P}(G) = P(G + uv) + P(G \setminus uv), \tag{7}$$

if $uv \notin E(G)$.

Since there is only one possible coloring for K_n (using exactly n colors), we have

$$P(K_n, k) = \begin{cases} 1 & \text{if } k = n, \\ 0 & \text{otherwise,} \end{cases}$$

and $\mathcal{P}(K_n) = 1$. This constitutes a base case for a straightforward recursive algorithm to compute $\mathcal{P}(G)$ for any graph G using relation (7). Another recursive procedure can be obtained from (6) using the empty graph $\overline{K_n}$ to define the base case. Indeed, we have

$$P(\overline{K_n}, k) = \begin{cases} \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} & \forall k \leq n, \\ 0 & \forall k > n, \end{cases}$$

where

$$\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = \frac{1}{k!} \sum_{j=1}^k (-1)^{k-j} \binom{k}{j} j^n$$

is a *Stirling number of the second kind*, that is the number of ways to partition a set of n elements into k non-empty subsets. It follows that

$$\mathcal{P}(\overline{K_n}) = \sum_{k=1}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = B_n,$$

where B_n is the n^{th} *Bell number* (sequence A000110 in OEIS [8]). This is not surprising since B_n represents the number of partitions of a set of n elements which is obviously the same as the number of non-equivalent colorings in a graph without any edge.

Of course, the complexities of the two above recursive algorithms are exponential in general. However, we will see in the next section that it can be refined to give a polynomial algorithm for some particular classes of graphs.

Generalized Stirling and Bell numbers have been defined and studied in [2] and are also linked to the new proposed invariant. More precisely, let

$$S_r(n, k) = \frac{1}{k!} \sum_{j=r}^k (-1)^{k-j} \binom{k}{j} \left(\frac{j!}{(j-r)!} \right)^n.$$

Consider n sets E_1, E_2, \dots, E_n of r elements. The *generalized Stirling number* $S_r(n, k)$ is the number of different partitions of these nr elements into k non-empty subsets such that each subset contains at most one element of each E_i . In other words, $S_r(n, k) = P(nK_r, k)$. The *Generalized Bell numbers* $B_{r,n}$ are then defined as follows:

$$B_{r,n} = \sum_{k=r}^{rn} S_r(n, k).$$

They represent the number of partitions of the nr elements so that each subset contains at most one element of each E_i . Hence, $B_{r,n} = \mathcal{P}(nK_r)$.

5 Determining some numbers of colorings

In this section, we determine the value of $\mathcal{P}(G)$ for several classes of graphs. We start with k -trees which are chordal graph with all maximal cliques of size $k + 1$ and all minimal clique separators of size k . Thus, a 1-tree is a tree. Note that every k -tree can be constructed from a complete graph on $k + 1$ vertices by adding vertices iteratively such that each new vertex has exactly k neighbors forming a clique [7]. To avoid confusion with the number of colors k we use the notation r -tree in the sequel.

Theorem 4 *Let T_n^r be a r -tree of order $n \geq r + 1$. Then,*

$$P(T_n^r, k) = \begin{Bmatrix} n - r \\ k - r \end{Bmatrix},$$

for all $k = r + 1, \dots, n$.

Proof. If $n = r + 1$, then T_n^r is a clique of size $r + 1$ and $P(T_n^r, n) = 1 = \begin{Bmatrix} n - r \\ n - r \end{Bmatrix}$. Otherwise, let v be a simplicial vertex of T_n^r . We consider two cases when counting the colorings of T_n^r : either v has its own color, or v has one color already used by other vertices of T_n^r . The first case gives $P(T_{n-1}^r, k - 1)$ colorings and the latter one $(k - r)P(T_{n-1}^r, k)$ colorings since the color of v cannot be the same than the r colors already used by its neighbors. Altogether we have

$$P(T_n^r, k) = P(T_{n-1}^r, k - 1) + (k - r)P(T_{n-1}^r, k).$$

Using induction, we get

$$P(T_n^r, k) = \begin{Bmatrix} n - 1 - r \\ k - 1 - r \end{Bmatrix} + (k - r) \begin{Bmatrix} n - 1 - r \\ k - r \end{Bmatrix} = \begin{Bmatrix} n - r \\ k - r \end{Bmatrix}.$$

The last equality comes from the known recurrence relations obeyed by Stirling numbers. □

Theorem 5 *Let T_n^r be a r -tree of order $n \geq r + 1$. Then,*

$$\mathcal{P}(T_n^r) = B_{n-r}.$$

Proof. By Theorem 4, and since any coloring of T_n^r has at least $r + 1$ colors,

$$\begin{aligned} \mathcal{P}(T_n^r) &= \sum_{k=r+1}^n P(T_n^r, k), \\ &= \sum_{k=r+1}^n \begin{Bmatrix} n - r \\ k - r \end{Bmatrix}, \\ &= \sum_{k=1}^{n-r} \begin{Bmatrix} n - r \\ k \end{Bmatrix} = B_{n-r}. \end{aligned}$$

□

Note that if $r = 0$, then r -trees are empty graphs and Theorem 5 is another way to show that $\mathcal{P}(\overline{K_n}) = B_n$. Another interesting particular case of r -trees are trees.

Corollary 6 *Let T be a tree of order $n \geq 1$. Then,*

$$\mathcal{P}(T) = B_{n-1}.$$

The decomposition used in the proof of Theorem 4 allows to compute the number of non-equivalent colorings of a chordal graph in polynomial time (using dynamic programming). Indeed, if v is a simplicial vertex of G with r neighbors, then

$$P(G, k) = P(G - v, k)(k - r) + P(G - v, k - 1).$$

Corollary 7 *If G is a chordal graph, then there exists a polynomial algorithm to compute $\mathcal{P}(G)$.*

Notice the same results also holds for the computation of the chromatic polynomial [6].

Theorem 8 *Let $K_{2,n}$ be a complete bipartite graph on $n + 2$ vertices such that $V = V_1 \cup V_2$ and $|V_1| = 2$. Then,*

$$\mathcal{P}(K_{2,n}) = 2B_n.$$

Proof. Let H be the graph obtained from $K_{2,n}$ with an additional edge between the two vertices of V_1 . Observe that H has a dominating vertex. Thus, applying (5), and then Property 1 and Corollary 6 gives

$$\mathcal{P}(K_{2,n}) = \mathcal{P}(K_{1,n}) + \mathcal{P}(H) = 2\mathcal{P}(K_{1,n}) = 2B_n.$$

□

Theorem 9 *Let C_n be a cycle of order $n \geq 3$. Then,*

$$\mathcal{P}(C_n) = \sum_{j=1}^{n-1} (-1)^{j+1} B_{n-j}.$$

Proof. Note that the result holds for $C_3 = K_3$ since $B_2 - B_1 = 1$. Applying (6) gives

$$\mathcal{P}(C_n) = \mathcal{P}(P_n) - \mathcal{P}(C_{n-1}).$$

Then, by Corollary 6 and the induction on n ,

$$\begin{aligned} \mathcal{P}(C_n) &= B_{n-1} - \sum_{j=1}^{n-2} (-1)^{j+1} B_{n-1-j}, \\ &= B_{n-1} - \sum_{j=2}^{n-1} (-1)^j B_{n-j}, \\ &= B_{n-1} + \sum_{j=2}^{n-1} (-1)^{j+1} B_{n-j}, \\ &= \sum_{j=1}^{n-1} (-1)^{j+1} B_{n-j}. \end{aligned}$$

□

Corollary 10 *Let W_n be a wheel of order $n \geq 4$. Then,*

$$\mathcal{P}(W_n) = \sum_{j=1}^{n-2} (-1)^{j+1} B_{n-j-1}.$$

Proof. By Property 1, we have $\mathcal{P}(W_n) = \mathcal{P}(C_{n-1})$, and the result follows from Theorem 9. □

Observe that Bell numbers appear repeatedly in the above results. Recall that B_n is the number of partitions of a set of n labeled elements without any constraint on the fact that two elements can be in the same partition or not. From a graph theoretical point of view, the partitions are the colors of the vertices and adding an edge represents such a constraint. In particular, it is of interest to note that the sequence $\mathcal{P}(C_n)$ for $n = 2, 3, \dots$ determined by Theorem 9 corresponds to sequence A000296 in OEIS [8]. This sequence is known to be the number of cyclically spaced partitions.

Given two graphs G and H of order n , we note $G >_P H$ and say that G *strictly dominates* H for the number of non-equivalent colorings if $P(G, k) \geq P(H, k)$ for all $k = 1, 2, \dots, n$, and there exists some integer k such that $P(G, k) > P(H, k)$. By Property 2, the following corollary is straightforward.

Corollary 11 *Let G, G' and H be three graphs such that G and G' have the same order. If $G >_P G'$, then, $\mathcal{P}(G \cup H) > \mathcal{P}(G' \cup H)$.*

6 Bounding the number of colorings of graphs with fixed maximum degree

In this section, we study upper and lower bounds on $\mathcal{P}(G)$ for graphs G with bounded maximum degree. We note that the following results were first conjectured with the help of the conjecture-making system GraPHedron [5].

The upper bound is straightforward. We define $G_{n,\Delta}^>$ to be the graph of order n and with a maximum degree Δ that is composed of a star $S_{\Delta+1}$ and $n - \Delta - 1$ isolated vertices (see Figure 5 for an example).




Figure 5: The graph $G_{8,4}^>$.

Theorem 12 *Let G be a graph of order n and maximum degree Δ . Then,*

$$\mathcal{P}(G) \leq \sum_{i=0}^{\Delta} (-1)^i \binom{\Delta}{i} \mathbf{B}_{n-i},$$

with equality if and only if G is isomorphic to $G_{n,\Delta}^>$.

Proof. The graph $G_{n,\Delta}^>$ is clearly the graph minimizing the number of edges among all graphs of order n with maximum degree Δ . Adding edges to $G_{n,\Delta}^>$ (in such a way that the maximum degree is not increased) will add new constraints between pairs of vertices, and this will therefore strictly decrease the number of colorings. Hence $\mathcal{P}(G) \leq \mathcal{P}(G_{n,\Delta}^>)$, with equality if and only if G is isomorphic to $G_{n,\Delta}^>$. It remains to prove that

$$\mathcal{P}(G_{n,\Delta}^>) = \sum_{i=0}^{\Delta} (-1)^i \binom{\Delta}{i} \mathbf{B}_{n-i} \quad \text{for all } n \text{ and } \Delta.$$

The equality holds for $\Delta = 0$ since $\mathcal{P}(G_{n,\Delta}^>)$ is then isomorphic to $\overline{K_n}$ and we have already observed that $\mathcal{P}(\overline{K_n}) = \mathbf{B}_n$. For larger values of Δ , we proceed by induction using the following equality obtained from (6):

$$\mathcal{P}(G_{n,\Delta}^>) = \mathcal{P}(G_{n,\Delta-1}^>) - \mathcal{P}(G_{n-1,\Delta-1}^>).$$

We then have

$$\begin{aligned} \mathcal{P}(G_{n,\Delta}^>) &= \sum_{i=0}^{\Delta-1} (-1)^i \binom{\Delta-1}{i} \mathbf{B}_{n-i} - \sum_{i=0}^{\Delta-1} (-1)^i \binom{\Delta-1}{i} \mathbf{B}_{n-1-i} \\ &= \sum_{i=0}^{\Delta-1} (-1)^i \binom{\Delta-1}{i} \mathbf{B}_{n-i} + \sum_{i=1}^{\Delta} (-1)^i \binom{\Delta-1}{i-1} \mathbf{B}_{n-i} \\ &= \mathbf{B}_n + \sum_{i=1}^{\Delta-1} (-1)^i \left(\binom{\Delta-1}{i} + \binom{\Delta-1}{i-1} \right) + (-1)^\Delta \mathbf{B}_{n-\Delta} \\ &= \sum_{i=0}^{\Delta} (-1)^i \binom{\Delta}{i} \mathbf{B}_{n-i}. \end{aligned}$$

□

A lower bound on $\mathcal{P}(G)$ for graphs of order n and bounded maximum degree Δ is easy to obtain for some values of Δ , but more intricate or still open for the other ones. In the rest of this section, we say that a graph G^* is *extremal* if $\mathcal{P}(G^*) \leq \mathcal{P}(G)$ for all graphs G of order n such that $\Delta(G) = \Delta(G^*)$. The following property will be used intensively in the ongoing proofs.

Property 13 *Let G be a graph with two vertices v and w such that $vw \notin E$ and*

$$\max(d(v), d(w)) < \Delta(G).$$

Then, G is not extremal.

Proof. Adding the edge vw will not change the value of $\Delta(G)$ but will strictly decrease the number of colorings of G . \square

We start by defining a graph of order n and with maximum degree equals to 1. If n is even, then $G_{n,\Delta=1}^<$ is the disjoint union of $\frac{n}{2}$ copies of K_2 ; if n is odd, it is the disjoint union of $G_{n-1,\Delta=1}^<$ and an isolated vertex. The graph $G_{7,\Delta=1}^<$ is drawn on the left-hand side of Figure 6.




Figure 6: The graphs $G_{7,\Delta=1}^<$, $G_{7,\Delta=2}^<$ and $K_6 \cup K_1$ (from left to right).

Theorem 14 Let G be a graph of order n such that $\Delta(G) = 1$. Then,

$$\mathcal{P}(G) \geq \sum_{i=0}^{\lfloor n/2 \rfloor} (-1)^i \binom{\lfloor n/2 \rfloor}{i} \mathcal{B}_{n-i},$$

with equality if and only if G is isomorphic to $G_{n,\Delta=1}^<$.

Proof. Since $\Delta(G) = 1$, G is a disjoint union of several copies of K_2 and isolated vertices. If G has at least two isolated vertices v and w , we know from Property 13 that it cannot be extremal. Thus, if G is extremal it must be isomorphic to $G_{n,\Delta=1}^<$.

Consider now the disjoint union of p K_2 and q K_1 . We prove that

$$\mathcal{P}(pK_2 \cup qK_1) = \sum_{i=0}^p (-1)^i \binom{p}{i} \mathcal{B}_{2p+q-i}.$$

The equality holds for $p = 0$ since the graph is then isomorphic to $\overline{K_q}$ and we have $\mathcal{P}(\overline{K_q}) = \mathcal{B}_q$. For larger values of p , we proceed by induction using the following equality obtained from (6):

$$\mathcal{P}(pK_2 \cup qK_1) = \mathcal{P}((p-1)K_2 \cup (q+2)K_1) - \mathcal{P}((p-1)K_2 \cup (q+1)K_1).$$

We then have

$$\begin{aligned} \mathcal{P}(pK_2 \cup qK_1) &= \sum_{i=0}^{p-1} (-1)^i \binom{p-1}{i} \mathcal{B}_{2p+q-i} - \sum_{i=0}^{p-1} (-1)^i \binom{p-1}{i} \mathcal{B}_{2p+q-1-i} \\ &= \sum_{i=0}^{p-1} (-1)^i \binom{p-1}{i} \mathcal{B}_{2p+q-i} + \sum_{i=1}^p (-1)^i \binom{p-1}{i-1} \mathcal{B}_{2p+q-i} \\ &= \mathcal{B}_{2p+q} + \sum_{i=1}^{p-1} (-1)^i \left(\binom{p-1}{i} + \binom{p-1}{i-1} \right) + (-1)^p \mathcal{B}_{p+q} \\ &= \sum_{i=0}^p (-1)^i \binom{p}{i} \mathcal{B}_{2p+q-i}. \end{aligned}$$

To conclude, it is sufficient to observe that $G_{n,\Delta=1}^<$ is isomorphic to $pK_2 \cup qK_1$ with $p = \lfloor n/2 \rfloor$ and $q = n - 2p$. \square

We now consider graphs G with maximum degree $\Delta(G) = 2$. Before giving a lower bound on $\mathcal{P}(G)$ for such graphs, we prove some useful lemmas.

Lemma 15 Consider a cycle C_n of order $n \geq 6$. Then,

$$\begin{aligned} \mathcal{P}(C_n, k) &> \mathcal{P}(C_{n-3} \cup C_3, k) \quad \text{for } k = 3, 4, \dots, n-2; \\ \mathcal{P}(C_n, k) &= \mathcal{P}(C_{n-3} \cup C_3, k) \quad \text{for } k = n-1, n. \end{aligned}$$

Proof. The values in the following table show that the result holds for $n = 6$.

k	3	4	5	6
$P(C_6, k)$	10	20	9	1
$P(2C_3, k)$	6	18	9	1

For larger values of n , the following equalities are obtained from (4) and (5):

$$\begin{aligned}
P(C_{n-3} \cup C_3, k) &= P(P_{n-3} \cup C_3, k) - P(C_{n-4} \cup C_3, k) \\
&= P(P_{n-3} \cup P_3, k) - P(P_{n-3} \cup P_2, k) - P(C_{n-4} \cup C_3, k) \\
&= (P(P_n, k) + P(P_{n-1}, k)) - (P(P_{n-1}, k) + P(P_{n-2}, k)) \\
&\quad - P(C_{n-4} \cup C_3, k) \\
&= P(P_n, k) - P(P_{n-2}, k) - P(C_{n-4} \cup C_3, k)
\end{aligned}$$

Clearly, $P(P_{n-2}, k) > 0$ for $k = 3, 4, \dots, n-2$ and $P(P_{n-2}, k) = 0$ for $k = n-1, n$. Also, by induction, we have $P(C_{n-4} \cup C_3, k) < P(C_{n-1}, k)$ for $k = 3, 4, \dots, n-2$, and $P(C_{n-4} \cup C_3, k) = P(C_{n-1}, k)$ for $k = n-1, n$. Hence, $P(C_{n-3} \cup C_3, k) \leq P(P_n, k) - P(C_{n-1}, k)$, with equality only if $k = n-1, n$. To conclude, we observe from (4) that $P(P_n, k) - P(C_{n-1}, k) = P(C_n, k)$. \square

Since $P(C_n, 2) \geq 0$ while $P(C_{n-3} \cup C_3, 2) = 0$ for $n \geq 6$, the following corollary is straightforward.

Corollary 16 Consider a cycle C_n of order $n \geq 6$. Then $C_n >_P C_{n-3} \cup C_3$.

Lemma 17 Consider a cycle C_n of order $n \geq 3$. Then,

$$P(C_n \cup K_1, k) = P(P_{n+1}, k) \quad \text{for } k = 3, 4, \dots, n+1.$$

Proof. The result is valid for $n = 3$ since $P(C_3 \cup K_1, 3) = P(P_4, 3) = 3$ and $P(C_3 \cup K_1, 4) = P(P_4, 4) = 1$. For larger values of n and $k \geq 3$, we proceed by induction and apply (4) and (5) to obtain:

$$\begin{aligned}
P(C_n \cup K_1, k) &= P(P_n \cup K_1, k) - P(C_{n-1} \cup K_1, k) \\
&= P(P_{n+1}, k) + P(P_n, k) - P(C_{n-1} \cup K_1, k) \\
&= P(P_{n+1}, k)
\end{aligned}$$

\square

Corollary 18 Consider a cycle C_n of order $n \geq 4$. Then

$$\begin{aligned}
C_n \cup K_1 &>_P C_{n+1} && \text{if } n \text{ is even;} \\
C_n \cup K_1 &>_P C_{n-2} \cup C_3 && \text{if } n \text{ is odd.}
\end{aligned}$$

Proof. Since $P(P_{n+1}, k) > P(C_{n+1}, k)$ for $k = 3, 4, \dots, n$, it follows from Lemma 17 that $P(C_n \cup K_1, k) > P(C_{n+1}, k)$ for $k = 3, 4, \dots, n$.

- If n is even, then $P(C_n \cup K_1, 2) = 2 > 0 = P(C_{n+1}, 2)$ and $P(C_n \cup K_1, n+1) = P(C_{n+1}, n+1) = 1$, which implies $C_n \cup K_1 >_P C_{n+1}$.
- If n is odd, then we know from Lemma 15 that $P(C_{n+1}, k) \geq P(C_{n-2} \cup C_3, k)$ for $k = 3, 4, \dots, n$. Since $P(C_n \cup K_1, 2) = P(C_{n-2} \cup C_3, 2) = 0$ and $P(C_n \cup K_1, n+1) = P(C_{n-2} \cup C_3, n+1) = 1$, we have $C_n \cup K_1 >_P C_{n-2} \cup C_3$.

\square

Lemma 19 Consider a cycle C_n of order $n \geq 5$. Then, $C_{n-2} \cup K_2 >_P C_n$.

Proof. By applying (4) and (5), we obtain the following equalities which are valid for all $k \geq 2$:

$$\begin{aligned}
P(C_{n-2} \cup K_2, k) &= P(P_{n-2} \cup K_2, k) - P(C_{n-3} \cup K_2, k) \\
&= P(P_n, k) + P(P_{n-1}, k) - P(P_{n-3} \cup K_2, k) \\
&\quad + P(C_{n-4} \cup K_2, k) \\
&= P(P_n, k) + P(P_{n-1}, k) - P(P_{n-1}, k) - P(P_{n-2}, k) \\
&\quad + P(C_{n-4} \cup K_2, k) \\
&= P(P_n, k) - P(P_{n-2}, k) + P(C_{n-4} \cup K_2, k).
\end{aligned} \tag{8}$$

We now analyse three different cases.

- If $k \geq 4$, we first show that $P(C_{n-2} \cup K_2, k) = P(P_n, k)$. This is true for $n = 5, 6$ since $P(C_3 \cup K_2, 4) = P(P_5, 4) = 6$, $P(C_3 \cup K_2, 5) = P(P_5, 5) = 1$, $P(C_4 \cup K_2, 4) = P(P_6, 4) = 25$, $P(C_4 \cup K_2, 5) = P(P_6, 5) = 10$, and $P(C_4 \cup K_2, 6) = P(P_6, 6) = 1$. For larger values of n , the equality is obtained by induction, using equation (8), since $P(C_{n-4} \cup K_2, k)$ is then equal to $P(P_{n-2}, k)$. Since $P(C_n, k) \leq P(P_n, k)$ for all k , we have $P(C_n, k) \leq P(C_{n-2} \cup K_2, k)$ for $k = 4, 5, \dots, n$.
- If $k = 3$, we first show that $P(C_{n-2} \cup K_2, 3) = P(P_n, 3) + (-1)^n$. This is true for $n = 5, 6$ since $P(C_3 \cup K_2, 3) = 6$, $P(P_5, 3) = 7$, $P(C_4 \cup K_2, 3) = 16$, and $P(P_6, 3) = 15$. For larger values of n , the equality is obtained by induction, using equation (8), since $P(C_{n-4} \cup K_2, k)$ is then equal to $P(P_{n-2}, 3) + (-1)^{n-2} = P(P_{n-2}, 3) + (-1)^n$. Since $P(C_{n-1}, 3) > 1$ for all $n \geq 5$, we conclude that $P(C_n, 3) = P(P_n, 3) - P(C_{n-1}, 3) \leq P(P_n, 3) - 2 < P(P_n, 3) + (-1)^n = P(C_{n-2} \cup K_2, 3)$.
- If $k = 2$ then $P(C_n, 2) \leq P(C_{n-2} \cup K_2, 2)$ since both $P(C_n, 2)$ and $P(C_{n-2} \cup K_2, 2)$ equal 0 if n is odd, while $P(C_n, 2) = 1 < 2 = P(C_{n-2} \cup K_2, 2)$ if n is even.

□

The graph $G_{n, \Delta=2}^<$ is defined as follows:

- it is the disjoint union of $\frac{n}{3}$ copies of K_3 if $n \equiv 0 \pmod{3}$;
- it is the disjoint union of $G_{n-4, \Delta=2}^<$ and C_4 if $n \equiv 1 \pmod{3}$;
- it is the disjoint union of $G_{n-5, \Delta=2}^<$ and C_5 if $n \equiv 2 \pmod{3}$.

The graph $G_{7, \Delta=2}^<$ is illustrated in the middle of Figure 6. We now give a lower bound on $\mathcal{P}(G)$ for graphs G with maximum degree $\Delta(G) = 2$ and order $n \geq 5$. This is not restrictive because if $n \leq 4$ and $\Delta = 2$, then $\Delta = n - 2$ or $\Delta = n - 1$ and these cases are treated later.

Theorem 20 *Let G be a graph of order $n \geq 5$ such that $\Delta(G) = 2$. Then,*

$$\mathcal{P}(G_{n, \Delta=2}^<) \leq \mathcal{P}(G),$$

with equality if and only if G is isomorphic to $G_{n, \Delta=2}^<$.

Proof. Suppose G is extremal. Since $\Delta(G) = 2$, G is the disjoint union of cycles and paths. It follows from Property 13 that at most one connected component of G is a path, and such a path can only be K_1 or K_2 .

Case 1: K_1 is a connected component of G .

Let C_r ($r \geq 3$) be a longest cycle of G . If $r = 3$, then G is the disjoint union of K_1 and at least two copies of C_3 (because $n \geq 5$). Thus, $G = 2C_3 \cup K_1 \cup H$ where H is a (possibly empty) disjoint union of C_3 . The following table shows that G is not extreme since $2C_3 \cup K_1 >_P C_3 \cup C_4$, a contradiction.

k	2	3	4	5	6	7
$P(K_1 \cup 2C_3, k)$	0	18	78	63	15	1
$P(C_3 \cup C_4, k)$	0	18	66	55	14	1

If $r \geq 4$, then we know from Corollary 18 that either C_{r+1} (if r is even) or $C_{r-2} \cup C_3$ (if r is odd) is strictly dominated by $C_r \cup K_1$. Hence, G is not extremal, a contradiction.

Case 2: K_2 is a connected component of G .

Let C_r be any cycle in G . We know from Lemma 19 that $C_r \cup K_2 >_P C_{r+2}$, which means that G is not extremal, a contradiction.

Case 3: G is the disjoint union of cycles.

Since G is extremal, we know from Corollary 16 that these cycles are copies of C_3 , C_4 or C_5 . The following tables show that $2C_5 >_P 2C_3 \cup C_4$, $C_5 \cup C_4 >_P 3C_3$, and $2C_4 >_P 2C_5 \cup C_3$. Hence, since G is extremal, it contains no more than one C_4 or one C_5 , which means that G is isomorphic to $G_{n,\Delta=2}^<$.

k	2	3	4	5	6	7	8	9	10
$P(2C_5, k)$	0	150	2250	6345	6025	2400	435	35	1
$P(2C_3 \cup C_4, k)$	0	108	1908	5838	5790	2361	433	35	1

k	2	3	4	5	6	7	8	9
$P(C_5 \cup C_4, k)$	0	90	750	1415	925	246	27	1
$P(3C_3, k)$	0	36	540	1242	882	243	27	1

k	2	3	4	5	6	7	8
$P(2C_4, k)$	2	52	241	296	126	20	1
$P(C_5 \cup C_3, k)$	0	30	210	285	125	20	1

□

Since $C_3 = K_3$, we can link the above result with the generalized Bell numbers mentioned in Section 4.

Corollary 21 *Let G be a graph of order n such that $n \equiv 0 \pmod{3}$ and $\Delta(G) = 2$. Then*

$$\mathcal{P}(G) \geq B_{3, \frac{n}{3}}$$

We now give a lower bound on $\mathcal{P}(G)$ for graphs G of order n and maximum degree $n - 2$.

Theorem 22 *Let G be a graph of order $n \geq 2$ such that $\Delta(G) = n - 2$. Then,*

$$\mathcal{P}(G) \geq n$$

with equality if and only if G is isomorphic to $K_{n-1} \cup K_1$ when $n \neq 4$, and G is isomorphic to $K_3 \cup K_1$ or C_4 otherwise.

Proof. The proof is by induction on n and the result is clearly valid for $n = 2$. Notice first that $\mathcal{P}(K_{n-1} \cup K_1) = n$ because either the isolated vertex of K_1 has its own color, or it uses one of the $n - 1$ colors in K_{n-1} . So let G be an extremal graph of order $n > 2$ with $\Delta(G) = n - 2$. We then have $\mathcal{P}(G) \leq \mathcal{P}(K_{n-1} \cup K_1) = n$. Let x be any vertex of degree $n - 2$, and let y be the unique vertex that is not adjacent to x . It follows from Property 13 that if two vertices v and w distinct from x and y are non-adjacent, then they are both adjacent to y . Hence, if y is an isolated vertex in G , then G is isomorphic to $K_{n-1} \cup K_1$.

So suppose $d(y) \geq 1$ and let v be one of its neighbors. Since v is not dominating, there exists at least one vertex w not adjacent to v . As observed above, w is necessarily adjacent to y . Let W be the set of vertices adjacent to y . We therefore have $|W| \geq 2$ and, by Property 13, every vertex non-adjacent to y has degree $n - 2$. Let G' be the graph induced by W . No vertex of G' is dominating (else it would also be dominating in G), and since at least one of v and w has degree $n - 2$ in G (and thus has degree $|W| - 2$ in G'), we have $\Delta(G') = |W| - 2$. By induction, $\mathcal{P}(G') \geq |W|$.

Given any coloring of G' , we can construct $n - |W|$ non-equivalent colorings of G by copying the colors on the vertices of W , assigning new colors to all vertices non-adjacent to y , and either assigning one of these $n - |W| - 1$ new colors to y , or a new one not shared by any other vertex. Hence,

$$n \geq \mathcal{P}(G) \geq \mathcal{P}(G')(n - |W|) \geq |W|(n - |W|). \quad (9)$$

Then, $n - |W| \geq |W|(n - |W| - 1) \geq 2(n - |W| - 1)$, which implies $n - |W| \leq 2$. Since x and y do not belong to W , we have $n - |W| = 2$. Hence, equation (9) becomes $|W| + 2 \geq 2|W|$, which is equivalent to $|W| \leq 2$. Since v and w belong to W , we have $|W| = 2$. In summary, $\mathcal{P}(G) = n = 4$ and G is isomorphic to C_4 . \square

Finally, notice that the lower bound on $\mathcal{P}(G)$ for graphs G with $\Delta(G) = n - 1$ is trivial since K_n has clearly the minimum number of colorings among all graph of order n .

7 Concluding remarks and open problems

We have defined a graph invariant that corresponds to the number of non-equivalent proper vertex colorings of a graph. We have shown similarities and differences between this invariant and the famous chromatic polynomial. We have also determined the value of this invariant for several classes of graphs and have given lower and upper bounds on its value for graphs with bounded maximum degree.

It would be interesting to determine a lower bound on $\mathcal{P}(G)$ for graphs G of order n and with maximum degree in $\{3, 4, \dots, n - 3\}$. The extremal graphs in this case do not seem to have a simple structure, as was the case for $\Delta(G) = 1, 2, n - 2, n$. We have determined some of them by exhaustive enumeration. For example, we have drawn in Figure 7 the only graphs G of order $n = 6, 7, 8, 9$ with minimum value $\mathcal{P}(G)$ when $\Delta(G) = 3, 4, 5$.

Notice also that several graphs with minimum value $\mathcal{P}(G)$ are non-connected. It would be interesting to determine these extremal graphs with the additional constraint that G must be connected. Also, it could be interesting to characterize the graphs G that minimize or maximize $\mathcal{P}(G)$ when the order and the size of G are fixed.













	$\Delta(G) = 3$	$\Delta(G) = 4$	$\Delta(G) = 5$
$n = 6$	 $\mathcal{P}(G) = 18$	 $\mathcal{P}(G) = 6$	 $\mathcal{P}(G) = 1$
$n = 7$	 $\mathcal{P}(G) = 70$	 $\mathcal{P}(G) = 29$	 $\mathcal{P}(G) = 7$
$n = 8$	 $\mathcal{P}(G) = 209$	 $\mathcal{P}(G) = 106$	 $\mathcal{P}(G) = 43$
$n = 9$	 $\mathcal{P}(G) = 1274$	 $\mathcal{P}(G) = 456$	 $\mathcal{P}(G) = 202$

Figure 7: Unique graphs G of order $n = 6, 7, 8, 9$ and maximum degree $\Delta(G) = 3, 4, 5$ with minimum value for $\mathcal{P}(G)$.

References

- [1] BIRKHOFF, G.D. A Determinant Formula for the Number of Ways of Coloring a Map. *Ann. Math. Harvard Coll.* 14 (1912), 42–46.
- [2] BLASIAK, P., PENSON, K.A., SOLOMON, A.I. The Boson Normal Ordering Problem and Generalized Bell Numbers *Annals of Combinatorics* 7 (2003), 127–139 .
- [3] DIESTEL, R. *Graph Theory*, second edition, Springer-Verlag, 2000.
- [4] DONG, F.M., KOH, K.M., TEO, K.L. *Chromatic Polynomials and Chromaticity of Graphs*, World Scientific Publishing Company, 2005, ISBN 981-256-317-2.
- [5] MÉLOT, H. Facet defining inequalities among graph invariants: the system GraPHedron. *Discrete Applied Mathematics* 156 (2008), 1875–1891.
- [6] NAOR, J., NAOR, M., SCHAFFER, A. Fast parallel algorithms for chordal graphs. *Proc. 19th ACM Symp. Theory of Computing* (1987), 355–364.
- [7] PATIL, H.P. On the structure of k -trees. *Journal of Combinatorics, Information and System Sciences* 11 (1986), 57–64.
- [8] SLOANE, N. The on-line encyclopedia of integer sequences. Available at <http://www.research.att.com/minjas/sequences/>.