

**The geometric–arithmetic index
and the chromatic number of
connected graphs**

M. Aouchiche
P. Hansen

G–2016–37

June 2016

Cette version est mise à votre disposition conformément à la politique de libre accès aux publications des organismes subventionnaires canadiens et québécois.

Avant de citer ce rapport, veuillez visiter notre site Web (<https://www.gerad.ca/fr/papers/G-2016-37>) afin de mettre à jour vos données de référence, s’il a été publié dans une revue scientifique.

This version is available to you under the open access policy of Canadian and Quebec funding agencies.

Before citing this report, please visit our website (<https://www.gerad.ca/en/papers/G-2016-37>) to update your reference data, if it has been published in a scientific journal.

Les textes publiés dans la série des rapports de recherche *Les Cahiers du GERAD* n’engagent que la responsabilité de leurs auteurs.

La publication de ces rapports de recherche est rendue possible grâce au soutien de HEC Montréal, Polytechnique Montréal, Université McGill, Université du Québec à Montréal, ainsi que du Fonds de recherche du Québec – Nature et technologies.

Dépôt légal – Bibliothèque et Archives nationales du Québec, 2016
– Bibliothèque et Archives Canada, 2016

The authors are exclusively responsible for the content of their research papers published in the series *Les Cahiers du GERAD*.

The publication of these research reports is made possible thanks to the support of HEC Montréal, Polytechnique Montréal, McGill University, Université du Québec à Montréal, as well as the Fonds de recherche du Québec – Nature et technologies.

Legal deposit – Bibliothèque et Archives nationales du Québec, 2016
– Library and Archives Canada, 2016

The geometric–arithmetic index and the chromatic number of connected graphs

Mustapha Aouchiche

Pierre Hansen

*GERAD & HEC Montréal, Montréal (Québec)
Canada, H3T 2A7*

mustapha.aouchiche@gerad.ca
pierre.hansen@gerad.ca

June 2016

**Les Cahiers du GERAD
G–2016–37**

Copyright © 2016 GERAD

Abstract: In the present paper, we compare the geometric–arithmetic index GA and the chromatic number χ of a connected graph with given order. We prove, among other results, an upper bound on the ratio GA/χ . We also prove lower bounds on the chromatic number in terms of geometric–arithmetic index and number of vertices of a connected graph. The results obtained for the chromatic number χ are extended to the clique number ω .

Keywords: Graph, geometric–arithmetic index, chromatic number, clique number, conjecture

Résumé: Dans le présent article, nous comparons l'indice géométrique-arithmétique GA et le nombre chromatique χ d'un graphe connexe d'ordre donné. Entre autres résultats, nous démontrons une borne supérieure sur le rapport GA/χ . Nous démontrons aussi des bornes inférieures sur le nombre chromatique en fonction de l'indice géométrique-arithmétique et du nombre de sommets d'un graphe connexe. Les résultats obtenus pour le nombre chromatique χ sont étendus au nombre de la clique maximum ω .

Mots clés: Graphe, indice géométrique–arithmtique, nombre chromatique, nombre de la clique maximum, conjecture

1 Introduction and definitions

We begin by recalling some definitions. In this paper, we consider only simple, undirected and finite graphs, *i.e.*, undirected graphs on a finite number of vertices without multiple edges or loops. A graph is (usually) denoted by $G = G(V, E)$, where V is its vertex set and E its edge set. The *order* of G is the number $n = |V|$ of its vertices and its *size* is the number $m = |E|$ of its edges.

As usual, we denote by P_n the path, by C_n the cycle, by S_n the star, by $K_{a,n-a}$ the complete bipartite graph and by K_n the complete graph, each on n vertices.

Molecular descriptors play a very important role in mathematical chemistry especially in QSAR (quantitative structure–activity relationship) and/or QSPR (quantitative structure–property relationship) related studies. Among those descriptors, a special interest is devoted to so-called topological indices. They are used to understand physicochemical properties of chemical compounds in a simple way, since they sum up some of the properties of a molecule in a single number. During the last decades, a legion of topological indices were introduced and found some applications in chemistry, see *e.g.*, [12, 13, 25]. The study of topological indices goes back to the seminal work by Wiener [28] in which he used the sum of all shortest-path distances, nowadays known as the *Wiener index*, of a (molecular) graph for modeling physical properties of alkanes.

Another very important molecular descriptor, was introduced by Randić [20]. It is called the *Randić (connectivity) index* and defined as

$$Ra = Ra(G) = \sum_{uv \in E} \frac{1}{\sqrt{d_u d_v}}$$

where d_u denotes the degree (number of neighbors) of u in G . The Randić index is probably the most studied molecular descriptor in mathematical chemistry. Actually, there are more than two thousand papers and five books devoted to this index (see, *e.g.*, [11, 15, 16, 18, 19] and the references therein).

Motivated by the definition of Randić connectivity index, Vukičević and Furtula [27] proposed the *geometric–arithmetic index*. It is so-called since its definition involves both the geometric and the arithmetic means of the endpoints degrees of the edges in a graph. For a simple graph G with edge set $E(G)$, the geometric–arithmetic index $GA(G)$ of a graph G is defined as in [27] by

$$GA(G) = \sum_{uv \in E(G)} \frac{2\sqrt{d_u d_v}}{d_u + d_v}.$$

where d_u denotes the degree of u in G .

It is noted in [27] that the predictive power of GA for physico-chemical properties is somewhat better than the predictive power of the Randić connectivity index. In [27], Vukičević and Furtula gave lower and upper bounds for GA , and identified the trees with minimum and maximum GA indices, which are the star and the path respectively. In [29] Yuan, Zhou and Trinajsić gave lower and upper bounds for GA index of molecular graphs using the numbers of vertices and edges. They also determined the n -vertex molecular trees with the minimum, the second, and the third minimum, as well as the second and third maximum GA indices. The chemical applicability of the geometric–arithmetic index was highlighted in [8, 10, 27].

Lower and upper bound on the geometric–arithmetic index in terms of order n , size m , minimum degree δ and/or maximum degree were proved in [21]. Also in [21], GA was compared to several other well known topological indices such as the Randić index, the first and second Zagreb indices, the harmonic index and the sum connectivity index. Other lower and upper bounds, on the geometric–arithmetic index, involving the order n the size m , the minimum and maximum degrees and the second Zagreb index were proved in [7].

In [1], several bounds and comparisons, involving the geometric–arithmetic index and several other graph parameters, were proved.

The problem of lower bounding GA over the class of connected graphs with fixed order of vertices and minimum degree was discussed in [9, 23].

Our main results are stated and proved in the next section. The third and last section is devoted to the statement of a few conjectures obtained after experiments using the computerized conjecture making system AutoGraphiX [2, 3, 5, 6].

2 Main results

A *coloring* of G is an assignment of colors to the vertices of G such that two adjacent vertices have different colors. The minimum number of colors in a coloring of G is the *chromatic number* of G and is denoted by $\chi(G)$. The chromatic number χ is a very widely studied graph invariant, whose history started with the famous *four color problem*, posed by Guthrie in 1852 (see e.g. [4, 22, 24] and the work of Kempe [17] in 1879 and Heawood [14] in 1890).

A *clique* of G is a subset of mutually adjacent vertices in G . A clique is called *maximal* if it is not contained in any other clique. A clique is called *maximum* if it has maximum cardinality. The maximum size of a clique in G is called the *clique number* of G and is denoted by $\omega = \omega(G)$.

In this section, we compare the geometric–arithmetic index GA and the chromatic number χ of a connected graph with given order. Results obtained for the chromatic number χ are extended to the clique number ω .

Note that all results proved in the present paper were first conjectured, or at least tested, using the conjecture–making system in graph theory AutoGraphix [2, 3, 5, 6].

We first prove an upper bound on the ratio GA/χ in terms of the number of vertices. We also characterize the corresponding extremal graphs.

Theorem 1 *For any connected graph on $n \geq 2$ vertices with chromatic number χ*

$$\frac{GA}{\chi} \leq \begin{cases} \frac{n^2}{8} & \text{if } n \text{ is even,} \\ \frac{(n^2-1)^{\frac{3}{2}}}{4n} & \text{if } n \text{ is odd,} \end{cases}$$

with equality if and only if G is the complete bipartite graph $K_{\frac{n}{2}, \frac{n}{2}}$ when n is even, and if and only if G is the complete bipartite graph $K_{\frac{n+1}{2}, \frac{n-1}{2}}$ when n is odd.

Proof. We have

$$GA(K_{\frac{n}{2}, \frac{n}{2}}) = \frac{n^2}{4} \quad \text{and} \quad GA(K_{\frac{n+1}{2}, \frac{n-1}{2}}) = \frac{(n^2-1)^{\frac{3}{2}}}{4n}.$$

If $\chi = 2$ and n is even, then

$$\frac{GA}{\chi} = \frac{GA}{2} \leq \frac{m}{2} \leq \frac{n^2}{8}$$

with equality if and only if G is the complete bipartite graph $K_{\frac{n}{2}, \frac{n}{2}}$.

If $G \cong K_{\frac{n+1}{2}, \frac{n-1}{2}}$, then

$$\frac{GA}{\chi} = \frac{(n^2-1)^{\frac{3}{2}}}{8n}.$$

If $\chi = 2$, n is odd and $G \not\cong K_{\frac{n+1}{2}, \frac{n-1}{2}}$, then

$$\frac{GA}{\chi} = \frac{GA}{2} \leq \frac{m}{2} \leq \frac{n^2-5}{8} < \frac{(n^2-1)^{\frac{3}{2}}}{8n}.$$

The last strict inequality can be proved using a few algebraic manipulations.

If $\chi \geq 3$, using the well-known Turan's theorem [26] on the maximum size of a graph with chromatic number χ , we have

$$\frac{GA}{\chi} \leq m \leq \frac{n}{2\chi} \left(n - \frac{n}{\chi} \right) = \frac{n^2(\chi-1)}{2\chi^2}.$$

The last expression is decreasing with respect to χ , so it reaches its maximum for $\chi = 3$. Therefore

$$\frac{GA}{\chi} \leq \frac{n^2}{9}.$$

Again, using a few algebraic manipulations, we get

$$\frac{GA}{\chi} \leq \frac{n^2}{9} < \begin{cases} \frac{n^2}{8} & \text{if } n \text{ is even,} \\ \frac{(n^2-1)^{\frac{3}{2}}}{4n} & \text{if } n \text{ is odd,} \end{cases}$$

for all $n \geq 3$. □

In a similar way, we can prove the following result involving the clique number instead of the chromatic number.

Theorem 2 *For any connected graph on $n \geq 2$ vertices with clique number ω*

$$\frac{GA}{\omega} \leq \begin{cases} \frac{n^2}{8} & \text{if } n \text{ is even,} \\ \frac{(n^2-1)^{\frac{3}{2}}}{4n} & \text{if } n \text{ is odd,} \end{cases}$$

with equality if and only if G is the complete bipartite graph $K_{\frac{n}{2}, \frac{n}{2}}$ when n is even, and if and only if G is the complete bipartite graph $K_{\frac{n+1}{2}, \frac{n-1}{2}}$ when n is odd.

We next prove a lower bound on the chromatic number χ over all connected graphs, using the number of vertices n , the geometric–arithmetic index GA and the maximum degree Δ . We also characterize the corresponding extremal graphs.

Proposition 1 *Let G be a connected graph with chromatic number χ , the geometric–arithmetic index GA and maximum degree Δ , then*

$$\chi \geq \frac{4GA}{n\Delta}$$

with equality if and only if n is even and G is a regular bipartite graph.

Proof. We have

$$n\Delta \geq 2m \text{ with equality if and only if } G \text{ is regular;}$$

$$\chi \geq 2 \text{ with equality if and only if } G \text{ is bipartite;}$$

$$m \geq GA \text{ with equality if and only if } G \text{ is regular.}$$

Thus $\chi n\Delta \geq 4GA$ with equality if and only if G regular bipartite. □

Following the same steps as in the above proof, with $n\bar{d} = 2m$ instead of $n\Delta \geq 2m$, we get the next proposition.

Proposition 2 *Let G be a connected graph with chromatic number χ , geometric–arithmetic index GA and average degree \bar{d} , then*

$$\chi \geq \frac{4GA}{n\bar{d}}$$

with equality if and only if n is even and G is a regular bipartite graph.

Considering the clique number ω instead of the chromatic number χ in the above two propositions, we can prove in the same way the following result.

Proposition 3 *Let G be a connected graph with clique number ω , geometric–arithmetic index GA and average degree \bar{d} , then*

$$\omega \geq \frac{4GA}{n\bar{d}} \geq \frac{4GA}{n\Delta}$$

with equality if and only if n is even and G is a regular bipartite graph.

3 Conjectures

In this section we list a few conjectures obtained using AutoGraphiX [2, 3, 5, 6].

In [1], an upper bound on the chromatic number χ , in terms of the geometric–arithmetic index GA and the minimum degree δ , was proved. The theorem is as follows.

Theorem 3 ([1]) *Let G be a connected graph with chromatic number χ , geometric–arithmetic index GA and minimum degree $\delta \geq 2$, then*

$$\chi \leq \frac{2GA}{\delta}$$

with equality if and only if G is the complete graph K_n .

Motivated by the above result, we conducted experiments with the help of AutoGraphiX in order to know if we can state a similar results with average degree \bar{d} and maximum degree Δ instead of minimum degree δ . In the case of average degree, the experiments led to the statement of the following conjecture.

Conjecture 1 *Let G be a connected graph with chromatic number χ , geometric–arithmetic index GA and average degree \bar{d} , then*

$$\chi \leq \frac{2GA}{\bar{d}}$$

with equality if and only if G is the complete graph K_n .

Note that a similar conjecture can be stated using the clique number ω instead of the chromatic number χ . It is as follows.

Conjecture 2 *Let G be a connected graph with clique number ω , geometric–arithmetic index GA and average degree \bar{d} , then*

$$\omega \leq \frac{2GA}{\bar{d}}$$

with equality if and only if G is the complete graph K_n .

In the case of maximum degree, we cannot state conjectures similar to the above ones. Indeed, there exist graphs with $\chi < 2GA/\Delta$ and $\omega < 2GA/\Delta$, such as the complete bipartite graph $K_{\frac{n}{2}, \frac{n}{2}}$ (for which $\chi = \omega = 2 < 2GA/\Delta = n$ for $n \geq 4$), and others with $\chi > 2GA/\Delta$ and $\omega > 2GA/\Delta$, such as the star S_n (for which $\chi = \omega = 2 > 2GA/\Delta = 4\sqrt{n-1}/n$ for $n \geq 3$).

In order to find a lower bound on the ratio GA/χ , our experiments led to the next conjecture. First, we need the following definition. A *pineapple* $PA_{n,k}$ is the graph obtained from a clique (a set of mutual adjacent vertices) on k vertices by attaching $n-k$ pendant edges to one of its vertices. The pineapple $PA_{10,6}$ is illustrated in Figure 1.




Figure 1: The pineapple $PA_{10,6}$.

Conjecture 3 Among connected graphs on n vertices with geometric–arithmetic index GA and chromatic number χ , GA/χ is minimum for the pineapple $PA_{n,\chi}$.

Note that for a pineapple $PA_{n,\chi}$ with $2 \leq \chi \leq n - 1$, we have

$$GA(PA_{n,\chi}) = \frac{(\chi - 1)(\chi - 2)}{2} + \frac{2(\chi - 1)\sqrt{(\chi - 1)(n - 1)}}{n + \chi - 2} + \frac{2(n - \chi)\sqrt{n - 1}}{n}$$

Similarly, we can state the following conjecture.

Conjecture 4 Among connected graphs on n vertices with geometric–arithmetic index GA and clique number ω , GA/ω is minimum for the pineapple $PA_{n,\omega}$.

Our attempts to find a lower bound on the geometric–arithmetic index GA in terms of the chromatic number χ , led to the following conjecture.

Conjecture 5 Let G be a connected graph with chromatic number χ and geometric–arithmetic index GA , then

$$GA \geq \frac{\chi(\chi - 1)}{2}$$

with equality if and only if G is the complete graph K_n .

Similarly, we can state the following conjecture.

Conjecture 6 Let G be a connected graph with clique number ω and geometric–arithmetic index GA , then

$$GA \geq \frac{\omega(\omega - 1)}{2}$$

with equality if and only if G is the complete graph K_n .

References

- [1] A. Ali, A. A. Bhatti and Z. Raza, Further inequalities between vertex-degree-based topological indices. arXiv preprint arXiv:1401.7511, 2014.
- [2] M. Aouchiche, J.-M. Bonnefoy, A. Fidahoussen, G. Caporossi, P. Hansen, L. Hiesse, J. Lacheré and A. Monhait, Variable Neighborhood Search for Extremal Graphs. 14. The AutoGraphiX 2 System. In L. Liberti and N. Maculan (editors), *Global Optimization: From Theory to Implementation*, Springer (2006) 281–310.
- [3] M. Aouchiche, G. Caporossi and P. Hansen, Variable Neighborhood Search for Extremal Graphs. 20. Automated Comparison of Graph Invariants MATCH Commun. Math. Comput. Chem. 58 (2007) 365–384.
- [4] N. L. Biggs, E. K. Lloyd and R. J. Wilson, *Graph Theory 1736–1936*. Oxford University Press, USA, 1986.
- [5] G. Caporossi and P. Hansen, Variable Neighborhood Search for Extremal Graphs. I. The AutoGraphiX System. *Discrete Math.* 212 (2000) 29–44.
- [6] G. Caporossi and P. Hansen, Variable Neighborhood Search for Extremal Graphs. 5. Three Ways to Automate Finding Conjectures. *Discrete Math.* 276 (2004) 81–94.
- [7] K. C. Das, I. Gutman, B. Furtula, On the first geometric–arithmetic index of graphs. *Discrete Applied Mathematics*, 159 (2011) 2030–2037.
- [8] K.C. Das, I. Gutman, B. Furtula, Survey on geometric–arithmetic indices of graphs. *MATCH Commun. Math. Comput. Chem.* 65 (2011) 595–644.
- [9] T. Divnić, M. Milivojević and Lj. Pavlović, Extremal graphs for the geometric–arithmetic index with given minimum degree. *Discrete Applied Mathematics* 162 (2014) 386–390.
- [10] B. Furtula, I. Gutman, Geometric–arithmetic Indices. In: I. Gutman, B. Furtula (Eds.), *Novel Molecular Structure Descriptors? Theory and Applications*, vol. I, Univ. Kragujevac, Kragujevac, 2010, pp. 137–172.
- [11] I. Gutman, B. Furtula (Eds.), *Recent Results in the Theory of Randić Index*. Univ. Kragujevac, Kragujevac, 2008.
- [12] I. Gutman, B. Furtula (Eds.), *Novel Molecular Structure Descriptors - Theory and Applications I*, Univ. Kragujevac, Kragujevac, 2010.
- [13] I. Gutman, B. Furtula (Eds.), *Novel Molecular Structure Descriptors - Theory and Applications II*, Univ. Kragujevac, Kragujevac, 2010.
- [14] P. J. Heawood, Map Color Theorems. *Quart. J. Math.* 24 (1890) 332–338
- [15] L. B. Keir, L. H. Hall, *Molecular connectivity in chemistry and drug research*. Academic Press, New York 1976.
- [16] L. B. Keir, L. H. Hall, *Molecular connectivity in structural-activity analysis*. Research study, Letchworth, England, 1986.
- [17] A. B. Kempe, On the Geographical Problem of Four Colors. *Amer. J. Math.* 2 (1879) 193–204.
- [18] X. Li, I. Gutman, *Mathematical Aspects of Randić Type Molecular Structure Descriptors*, Univ. Kragujevac, Kragujevac, 2006.
- [19] X. Li, Y. Shi, A survey on the Randić index, *MATCH Commun. Math. Comput. Chem.* 59 (2008) 127–156.
- [20] M. Randić, On characterization of molecular branching. *J. Am. Chem. Soc.* 97 (1975) 6609–6615.
- [21] J. M. Rodríguez and J. M. Sigarreta, On the Geometric–Arithmetic Index. *MATCH Commun. Math. Comput. Chem.* 74 (2015) 103–120.
- [22] T. L. Saaty and P. C. Kainen, *Four–Color Problem*. Dover Publications, New York, 1986.
- [23] M. Sohrabi-Haghighat and M. Rostami, Using linear programming to find the extremal graphs with minimum degree 1 with respect to Geometric–Arithmetic index. *Applied mathematics in Engineering, Management and Technology* 3 (2015) 534–539.
- [24] A. Soifer, *The Mathematical Coloring Book: mathematics of coloring and the colorful life of its creators*. Springer, 2008.
- [25] R. Todeschini, V. Consonni, *Molecular Descriptors for Chemoinformatics*, Wiley-VCH, Weinheim, 2009.
- [26] P. Turán, Eine Extremalaufgabe aus der Graphentheorie. (Hungarian) *Mat. Fiz. Lapok* 48 (1941) 436–452.
- [27] D. Vukičević and B. Furtula, Topological index based on the ratios of geometrical and arithmetical means of end vertex degrees of edges. *Journal of Mathematical Chemistry* 46 (2009) 1369–1376.
- [28] H. Wiener, Structural determination of paraffin boiling points, *J. Am. Chem. Soc.* 69 (1947) 17–20.
- [29] Y. Yuan, B. Zhou, and N. Trinajstić, On geometric–arithmetic index. *Journal of Mathematical Chemistry* 47 (2010) 833–841.