

Risk-sensitive linear-quadratic-Gaussian graphon mean-field games

T. Chen, M. Huang

G-2025-57

August 2025

La collection *Les Cahiers du GERAD* est constituée des travaux de recherche menés par nos membres. La plupart de ces documents de travail a été soumis à des revues avec comité de révision. Lorsqu'un document est accepté et publié, le pdf original est retiré si c'est nécessaire et un lien vers l'article publié est ajouté.

Citation suggérée : T. Chen, M. Huang (Août 2025). Risk-sensitive linear-quadratic-Gaussian graphon mean-field games, Rapport technique, Les Cahiers du GERAD G- 2025-57, GERAD, HEC Montréal, Canada.

Avant de citer ce rapport technique, veuillez visiter notre site Web (<https://www.gerad.ca/fr/papers/G-2025-57>) afin de mettre à jour vos données de référence, s'il a été publié dans une revue scientifique.

The series *Les Cahiers du GERAD* consists of working papers carried out by our members. Most of these pre-prints have been submitted to peer-reviewed journals. When accepted and published, if necessary, the original pdf is removed and a link to the published article is added.

Suggested citation: T. Chen, M. Huang (August 2025). Risk-sensitive linear-quadratic-Gaussian graphon mean-field games, Technical report, Les Cahiers du GERAD G-2025-57, GERAD, HEC Montréal, Canada.

Before citing this technical report, please visit our website (<https://www.gerad.ca/en/papers/G-2025-57>) to update your reference data, if it has been published in a scientific journal.

La publication de ces rapports de recherche est rendue possible grâce au soutien de HEC Montréal, Polytechnique Montréal, Université McGill, Université du Québec à Montréal, ainsi que du Fonds de recherche du Québec – Nature et technologies.

Dépôt légal – Bibliothèque et Archives nationales du Québec, 2025
– Bibliothèque et Archives Canada, 2025

The publication of these research reports is made possible thanks to the support of HEC Montréal, Polytechnique Montréal, McGill University, Université du Québec à Montréal, as well as the Fonds de recherche du Québec – Nature et technologies.

Legal deposit – Bibliothèque et Archives nationales du Québec, 2025
– Library and Archives Canada, 2025

GERAD HEC Montréal
3000, chemin de la Côte-Sainte-Catherine
Montréal (Québec) Canada H3T 2A7

Tél. : 514 340-6053
Télec. : 514 340-5665
info@gerad.ca
www.gerad.ca

Risk-sensitive linear-quadratic-Gaussian graphon mean-field games

Tian Chen ^{a, b}

Minyi Huang ^{b, c}

^a School of Mathematics, Shandong University, Jinan, China, 250100

^b School of Mathematics and Statistics, Carleton University, Ottawa (On), Canada, K1S 5B6

^c GERAD, Montréal (Qc), Canada, H3T 1J4

chentian43@sdu.edu.cn

mhuang@math.carleton.ca

August 2025
Les Cahiers du GERAD
G–2025–57

Copyright © 2025 Chen, Huang

Les textes publiés dans la série des rapports de recherche *Les Cahiers du GERAD* n'engagent que la responsabilité de leurs auteurs. Les auteurs conservent leur droit d'auteur et leurs droits moraux sur leurs publications et les utilisateurs s'engagent à reconnaître et respecter les exigences légales associées à ces droits. Ainsi, les utilisateurs:

- Peuvent télécharger et imprimer une copie de toute publication du portail public aux fins d'étude ou de recherche privée;
- Ne peuvent pas distribuer le matériel ou l'utiliser pour une activité à but lucratif ou pour un gain commercial;
- Peuvent distribuer gratuitement l'URL identifiant la publication.

Si vous pensez que ce document enfreint le droit d'auteur, contactez-nous en fournissant des détails. Nous supprimerons immédiatement l'accès au travail et enquêterons sur votre demande.

The authors are exclusively responsible for the content of their research papers published in the series *Les Cahiers du GERAD*. Copyright and moral rights for the publications are retained by the authors and the users must commit themselves to recognize and abide the legal requirements associated with these rights. Thus, users:

- May download and print one copy of any publication from the public portal for the purpose of private study or research;
- May not further distribute the material or use it for any profit-making activity or commercial gain;
- May freely distribute the URL identifying the publication.

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract : This paper investigates a class of linear-quadratic-Gaussian risk-sensitive graphon mean-field games, involving an asymptotically infinite population of heterogeneous agents distributed across an asymptotically infinite network, where each agent aims to minimize an exponential cost functional reflecting its risk sensitivity. Following the Nash certainty equivalence methodology, an auxiliary risk-sensitive optimal control problem is constructed and further combined with a consistency condition to determine decentralized strategies of the agents. The well-posedness of the resulting graphon mean-field game equation system, consisting of a family of fully coupled forward-backward differential equations, is established by a fixed point approach under a contraction condition, and by the method of continuity under an operator monotonicity condition, respectively. To prove the ε -Nash equilibrium property of the obtained decentralized strategies, one faces significant challenge since the usual L^2 error estimates on mean-field approximations are no longer adequate due to unboundedness of the integrand in the exponentiated cost. The proof will be accomplished by establishing certain exponentiated error estimates instead of L^2 error estimates. Finally, a numerical example is provided to illustrate our results.

Keywords: Mean-field game, graphon, risk-sensitive, ε -Nash equilibrium

Acknowledgements: This work was supported by the National Natural Science Foundation of China under contract no. 12401583, and by Natural Sciences and Engineering Research Council of Canada.

1 Introduction

Mean-field game (MFG) theory has rapidly developed into a robust framework for addressing non-cooperative dynamic decision-making in large populations of comparably small agents [26, 27, 31]. Most existing analyses are based on two approaches: the direct approach and the fixed point approach. The direct approach solves an N -player game, resulting in a large-scale coupled equation system. As the number of agents $N \rightarrow \infty$, this system reduces to a Hamilton–Jacobi–Bellman (HJB) equation coupled with a Fokker–Planck–Kolmogorov (FPK) equation [11, 30, 31]. Alternatively, the fixed point approach starts by assuming a known mean-field effect and derives the best response control law of a representative agent. Subsequently, the above mean-field is required to be regenerated by the closed-loop dynamics of an infinite population. This solution procedure is formalized as a fixed point problem [7, 13, 26], and is designated as the Nash certainty equivalence methodology in [27]. Recently, [28] analyzed the exact relationship between these two approaches in the setup of linear-quadratic-Gaussian (LQG) MFGs. MFG theory has found wide applications in areas such as economics and finance [18, 22], production output adjustment [44], and traffic management [4], among others.

Graphon mean-field games (GMFGs) as introduced in [10] significantly extend the classical MFG framework by incorporating agents distributed across large networks. As networks increase in size and complexity, graphon theory becomes a powerful tool to study their limits, enabling analysis of dense networks [8, 9, 34, 35]. Graphons, as measurable functions representing the limit of increasingly large graphs, provide a rigorous mathematical framework for studying the behavior of large-scale networks. This framework is particularly studied for fields such as economics, control theory, social networks, and large-scale interacting particle systems, where agents demonstrate heterogeneous and complex interactions [2, 5, 12, 16, 21, 23]. GMFGs enable researchers to study the limiting behavior of systems with increasing populations of agents, capturing both local and global network interaction effects.

The aforementioned literature on MFGs or GMFGs has primarily focused on risk-neutral cost functionals, often neglecting agents’ attitudes towards risk. However, in many practical scenarios, risk sensitivity plays a crucial role in decision-making, as recognized very early in [24]. Jacobson [29] introduced a finite horizon risk-sensitive optimal control problem in an LQG setting, where the agent’s cost is specified as an exponential function of the accumulated cost over time. Since the early work in [24, 29], this class of control problems has been extensively studied in the literature; see [6, 17, 19, 20, 32, 38, 46]. Within the framework of MFGs, risk-sensitive costs were adopted in [43], which specified the mean-field equilibrium strategies using a corresponding pair of HJB and FPK equations. Subsequently, Moon and Başar [37] extended this line of research by employing the maximum principle to analyze nonlinear risk-sensitive MFG models. risk-sensitive MFGs with major-minor agents were studied in [14, 33], addressing the interaction structure between the major and minor agents. In the setting of discrete-time Markov decision processes, related results can be found in [41]. Recently, Wang and Huang [45] considered LQG risk-sensitive MFGs and proved an $O(1/N)$ -Nash equilibrium theorem by applying a re-scaling technique to a family of coupled-Riccati equation systems with increasing dimensions.

In this paper, we formulate a class of linear-quadratic-Gaussian GMFGs with risk-sensitive cost functionals. Using the Nash certainty equivalence methodology [26, 27], we solve the risk-sensitive best response control problem of a representative agent after approximating its graphon coupling term by a deterministic function, and subsequently impose a consistency condition on the proposed limiting function. This solution procedure leads to the GMFG equation system, as a family of fully coupled forward-backward equations, whose well-posedness is established via a fixed point approach under a contraction condition, and by the method of continuity under an operator monotonicity condition, respectively. In contrast to similar conditions in the literature (see [25, 40]), our monotonicity condition involves operators acting on an infinite dimensional space. Furthermore, by a novel method of exponentiated error estimate, we prove that the set of decentralized strategies computed via the GMFG equation system constitutes an ε -Nash equilibrium for the finite population of agents. When the running cost is unbounded as in the LQG case, one encounters significant difficulty proving that

the set of decentralized strategies obtained from the risk-sensitive MFG is an ε -Nash equilibrium. In this case, the usual L^2 error estimates for mean-field approximations is inadequate for handling exponentiated cost functionals; see Remark 5.7 later on. Moreover, the method of performance estimates in [45] is not applicable to the GMFGs due to lack of symmetry of the agents. To establish the ε -Nash equilibrium theorem, we take a different route by deriving new approximation error estimates of exponential type without imposing any additional restrictions as used in the literature (see discussions in [33, Section 4.4]).

The main contributions of this paper are summarized as follows:

- We introduce a novel risk-sensitive GMFG model, where the large population of agents is distributed over dense networks.
- The solution of GMFG is characterized by a fully coupled forward-backward equation system parameterized by the nodal index. The unique solvability of the GMFG equation system is analyzed by two methods: (i) the fixed point method under a contraction condition, (ii) the method of continuity under a new operator monotonicity condition.
- Based on the GMFG equation system, we construct decentralized feedback strategies in the N -player model. A novel method via exponentiated error estimates is applied to prove that the set of decentralized strategies constitutes an ε -Nash equilibrium.

The paper is organized as follows. Section 2 formulates the risk-sensitive GMFG. Section 3 solves the best response control problem in a graphon mean-field limit model. By imposing a consistency condition, Section 4 derives the GMFG equation system and investigates its solvability by two approaches: the fixed point method and the method of continuity. In Section 5, we prove an ε -Nash equilibrium theorem for the obtained decentralized strategies. Section 6 presents a numerical example. Section 7 concludes the paper.

2 Problem formulation and preliminary

We use \mathbb{R}^m to denote the m -dimensional Euclidean space, consisting of column vectors, with norm $|\cdot|$ and inner product $\langle \cdot, \cdot \rangle$. Let D^\top (resp., D^{-1}) stand for the transpose (resp., inverse) of a matrix D , and \mathbb{S}^m for the set of symmetric real $m \times m$ matrices. If $D \in \mathbb{S}^n$ is positive definite (resp., positive semi-definite), we write $D > 0$ (resp., $D \geq 0$). Moreover, if an \mathbb{S}^n -valued deterministic function D defined on $[0, T]$ is uniformly positive definite, we write $D \gg 0$. For two sequences of number $\{a_k\}_{k \geq 1}$ and $\{b_k\}_{k \geq 1}$ with $b_k > 0$, if $\lim_{k \rightarrow \infty} \frac{a_k}{b_k} = 0$, we write $a_k = o(b_k)$.

Let $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ be a complete filtered probability space with filtration $\mathbb{F} := \{\mathcal{F}_t\}_{0 \leq t \leq T}$. Given Hilbert space \mathbb{H} and interval $[0, T]$, let $L^\infty([0, T]; \mathbb{H})$ stand for the space of all \mathbb{H} -valued essentially bounded functions defined on $[0, T]$, $C([0, T]; \mathbb{H})$ for the space of all \mathbb{H} -valued continuous functions, $L^2_{\mathbb{F}}([0, T]; \mathbb{H})$ for the space of all \mathbb{H} -valued, \mathcal{F}_t -adapted, square-integrable processes, and $S^2_{\mathbb{F}}([0, T]; \mathbb{H})$ for the space of all \mathbb{H} -valued, \mathcal{F}_t -adapted continuous processes ϕ such that $\mathbb{E}[\sup_{0 \leq t \leq T} |\phi(t)|^2] < \infty$.

2.1 Preliminary

The basic idea of the theory of graphons is that the edge structure of each finite cardinality network is represented by a step function density on the unit square in \mathbb{R}^2 on which the cut norm and cut metric (see [34]) are introduced. The set of finite graphs endowed with the cut metric then gives rise to a metric space, and the completion is the space of graphons. Let \mathbf{G}_0^{sp} be the linear space of bound symmetric Lebesgue measurable functions $W : [0, 1]^2 \rightarrow \mathbb{R}$, which are called kernels. The space \mathbf{G}^{sp} of graphons is a subset of \mathbf{G}_0^{sp} and consists of kernels $W : [0, 1]^2 \rightarrow [0, 1]$, which can be interpreted as weighted graphs on the vertex set $[0, 1]$.

In this paper, we start the modeling of the game of a finite population based on a finite graph. Specifically, the population resides on a weighted finite graph G_N with a set of nodes $\mathcal{V}_N = \{1, 2, \dots, N\}$

and weights $g_{ij}^N \in [0, 1]$ for $(i, j) \in \mathcal{V}_N \times \mathcal{V}_N$, where a value g_{ii}^N is still assigned in the case $i = j$. We call $g_i^N = (g_{i1}^N, g_{i2}^N, \dots, g_{iN}^N)$ a section of g^N at i . Moreover, each node l is occupied by exactly one agent, denoted by \mathcal{A}_l . In fact, it brings about no essential difficulty to consider the case where each node l is occupied by a subpopulation. Our further analysis is based on the convergence of g^N to a graphon limit g . We may naturally identify $(g_{ij}^N)_{1 \leq i, j \leq N}$ with a graphon $g^N(\alpha, \beta)$ as a step function defined on $[0, 1] \times [0, 1]$ (see [34]). We define the section of g at α by $g_\alpha : \beta \mapsto g(\alpha, \beta)$, $\beta \in [0, 1]$. We partition $[0, 1]$ into N subintervals of equal length. Here $I_1^N = [0, 1/N]$, $I_l^N = ((l-1)/N, l/N]$ for $2 \leq l \leq N$. When it is clear from the context, we omit the superscript N and write I_l . To relate the agents to the graphon vertex set $[0, 1]$, we let the i -th agent correspond to I_i .

2.2 Graphon MFG

For the risk-sensitive GMFG, consider a population of N agents $\{\mathcal{A}_i\}_{1 \leq i \leq N}$. The dynamics of agent \mathcal{A}_i residing at node i are governed by the following SDE:

$$dx_i(t) = [A(t)x_i(t) + B(t)u_i(t) + D(t)x_i^{(N)}(t)]dt + \sigma(t)dw_i(t), \quad x_i(0) = \xi_i, \quad (1)$$

where $x_i(t) \in \mathbb{R}^n$ is the state, $u_i(t) \in \mathbb{R}^m$ is the control, and $w_i, 1 \leq i \leq N$, are independent d -dimensional standard Brownian motions. Here, $x_i^{(N)} = \frac{1}{N} \sum_{j=1}^N g_{ij}^N x_j$ represents the weighted state-average of all agents with respect to node i .

Now we introduce two types of state feedback strategies. Let the functions $\varphi(t, x_1, \dots, x_N) : [0, T] \times \mathbb{R}^n \times \dots \times \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $\psi(t, x) : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^m$ be continuous in t , and Lipschitz continuous in (x_1, \dots, x_N) and x , respectively. For agent \mathcal{A}_i , we introduce

$$\mathcal{U}_c^i = \{u_i \mid u_i = \varphi(t, x_1, \dots, x_N)\}, \quad \mathcal{U}_d^i = \{u_i \mid u_i = \psi(t, x_i)\} \quad (2)$$

as the set of centralized (state) feedback strategies and the set of decentralized (state) feedback strategies, respectively. For simplicity, let $u = (u_1, \dots, u_N)$ be the set of strategies of N agents and $u_{-i} = (u_1, \dots, u_{i-1}, u_{i+1}, \dots, u_N)$ be the set of strategies of all agents other than \mathcal{A}_i . Note that while the coefficients are dependent on the time variable t , in further analysis, the variable t will usually be suppressed if no confusion occurs.

The risk-sensitive cost functional of agent \mathcal{A}_i takes the form

$$\mathcal{J}_i(u_i, u_{-i}) = \mathbb{E} \left[\exp \left(\gamma \left\{ \int_0^T [\|x_i - \Gamma x_i^{(N)}\|_Q^2 + \|u_i\|_R^2] dt + \|x_i(T) - \Gamma_f x_i^{(N)}(T)\|_{Q_f}^2 \right\} \right) \right], \quad (3)$$

where we denote $\|x\|_Q^2 = \langle Qx, x \rangle$, and both Q and R are matrix functions of t . Here $\gamma > 0$ is a given constant representing the risk sensitivity parameter.

For convenience of describing the asymptotic behavior of the agents and the networks, we need to relate the node of each agent to the continuum nodal set $[0, 1]$. Without loss of generality, we associate agent \mathcal{A}_i , $1 \leq i \leq N$, with the subinterval I_i in the partition of $[0, 1]$, which in turn determines the corresponding step function $g^N(\cdot, \cdot)$. We introduce the following assumptions.

- (H1) The initial states $\xi_i, 1 \leq i \leq N$, are independent random variables taking values in a compact set $S_0^x \subset \mathbb{R}^n$, and are independent of $\{w_i, 1 \leq i \leq N\}$. There exists a continuous function m_α^x defined on $[0, 1] \ni \alpha$, such that $\lim_{N \rightarrow \infty} \sup_{0 \leq \alpha \leq 1} |\mathbb{E} \xi_\alpha^N - m_\alpha^x| = 0$, where $\xi_\alpha^N = \sum_{i=1}^N \mathbb{1}_{I_i}(\alpha) \xi_i$.
- (H2) $A, D \in L^\infty([0, T]; \mathbb{R}^{n \times n})$, $B \in L^\infty([0, T]; \mathbb{R}^{n \times m})$, $\sigma \in L^\infty([0, T]; \mathbb{R}^{n \times d})$.
- (H3) $Q \in L^\infty([0, T]; \mathbb{S}^n)$, $R \in L^\infty([0, T]; \mathbb{S}^m)$, $Q_f \in \mathbb{S}^n$, $\Gamma, \Gamma_f \in \mathbb{R}^{n \times n}$, and $Q \geq 0$, $R \gg 0$, $Q_f \geq 0$.
- (H4) $BR^{-1}B^\top - 2\gamma\sigma\sigma^\top \geq 0$ for all $t \in [0, T]$.
- (H5) For any bounded, measurable function $h(\beta)$, the function $\int_0^1 g(\alpha, \beta)h(\beta)d\beta$ is continuous in $\alpha \in [0, 1]$.

Under (H1)–(H3), one can easily check that SDE (1) admits a unique solution $x_i \in S_{\mathbb{F}}^2([0, T]; \mathbb{R}^n)$ for any control $u_i \in L_{\mathbb{F}}^2([0, T]; \mathbb{R}^m)$ and the cost functional (3) is well-defined.

A basic solution notion for (1)–(3) is a Nash equilibrium (u_1^*, \dots, u_N^*) , where each u_i^* belongs to \mathcal{U}_c^i , corresponding to closed-loop perfect state (CLPS) information [3]. However, such a solution with its associated information pattern is impractical when the system consists of a large number of heterogeneous agents.

3 The design of decentralized strategies

To design decentralized strategies, we need to study an auxiliary best response control problem with an infinite population and a graphon limit g as an approximation of g^N (the nature of the approximation will be made exact by (H7) in Section 5). For an α -agent \mathcal{A}_α situated at vertex α , denote the graphon weighted mean state by $z_\alpha \in C([0, T]; \mathbb{R}^n)$, which is intended to approximate $x_i^{(N)}$ when the nodal location of \mathcal{A}_i is approximated by α . Let w_α be the standard Brownian motion associated with the α -agent. Then the α -agent has dynamics

$$dx_\alpha = (Ax_\alpha + Bu_\alpha + Dz_\alpha)dt + \sigma dw_\alpha, \quad x_\alpha(0) = \xi_\alpha, \quad \alpha \in [0, 1], \quad (4)$$

and cost functional

$$J_\alpha(u_\alpha) = \mathbb{E} \left[\exp \left(\gamma \left\{ \int_0^T [\|x_\alpha - \Gamma z_\alpha\|_Q^2 + \|u_\alpha\|_R^2] dt + \|x_\alpha(T) - \Gamma_f z_\alpha(T)\|_{Q_f}^2 \right\} \right) \right]. \quad (5)$$

The Brownian motion w_α and the initial state ξ_α are independent. The auxiliary best response (BR) control problem is formulated as follows.

Problem (BR). For the α -agent \mathcal{A}_α in (4)–(5), find a feedback control law \bar{u}_α such that

$$J_\alpha(\bar{u}_\alpha) = \inf_{u_\alpha} J_\alpha(u_\alpha).$$

For any given $t \in [0, T]$ and a deterministic initial condition $x_\alpha(t) = \mathbf{x} \in \mathbb{R}^n$, define

$$J_\alpha(t, \mathbf{x}; u_\alpha) = \mathbb{E} \left[\exp \left(\gamma \left\{ \int_t^T [\|x_\alpha - \Gamma z_\alpha\|_Q^2 + \|u_\alpha\|_R^2] ds + \|x_\alpha(T) - \Gamma_f z_\alpha(T)\|_{Q_f}^2 \right\} \right) \right].$$

and write the value function in the form

$$\inf_{u_\alpha} J_\alpha(t, \mathbf{x}; u_\alpha) = e^{\gamma V^\alpha(t, \mathbf{x})},$$

where the function V^α is to be determined. We write the gradient of a function as a row vector. Applying dynamic programming (see [6, Eqn. (1.4)]), we have

$$\begin{cases} -\frac{\partial V^\alpha(t, \mathbf{x})}{\partial t} = \inf_{u \in \mathbb{R}^m} \left\{ \frac{\partial V^\alpha(t, \mathbf{x})}{\partial \mathbf{x}} [A\mathbf{x} + Bu + Dz_\alpha(t)] + \|\mathbf{x} - \Gamma z_\alpha(t)\|_Q^2 + \|u\|_R^2 \right\} \\ \quad + \frac{1}{2} \text{Tr} \left(\sigma \sigma^\top \frac{\partial^2 V^\alpha(t, \mathbf{x})}{\partial \mathbf{x}^2} \right) + \frac{\gamma}{2} \left| \sigma^\top \left(\frac{\partial V^\alpha(t, \mathbf{x})}{\partial \mathbf{x}} \right)^\top \right|^2, \\ V^\alpha(T, \mathbf{x}) = \|\mathbf{x} - \Gamma_f z_\alpha(T)\|_{Q_f}^2, \quad (t, \mathbf{x}) \in [0, T] \times \mathbb{R}^n, \quad \alpha \in [0, 1], \end{cases} \quad (6)$$

and the optimal control law is given by

$$\bar{u}_\alpha(t) = -\frac{1}{2} R^{-1}(t) B^\top(t) \left(\frac{\partial V^\alpha(t, \mathbf{x})}{\partial \mathbf{x}} \right)^\top.$$

We write $V^\alpha(t, \mathbf{x}) = \mathbf{x}^\top \Pi(t) \mathbf{x} + 2\mathbf{x}^\top S_\alpha(t) + r_\alpha(t)$. Then by (6), we derive the following Riccati equation

$$\dot{\Pi} + \Pi A + A^\top \Pi - \Pi(BR^{-1}B^\top - 2\gamma\sigma\sigma^\top)\Pi + Q = 0, \quad \Pi(T) = Q_f. \quad (7)$$

Under (H2)–(H4), Riccati equation (7) admits a unique solution $\Pi \in C([0, T]; \mathbb{S}^n)$; see [1, pp. 186–190]. By (6), we further derive the following ordinary differential equations (ODEs):

$$\dot{S}_\alpha + (A^\top - \Pi BR^{-1}B^\top + 2\gamma\Pi\sigma\sigma^\top)S_\alpha - (Q\Gamma - \Pi D)z_\alpha = 0, \quad (8)$$

where $S_\alpha(T) = -Q_f\Gamma_f z_\alpha(T)$, and

$$\dot{r}_\alpha - S_\alpha^\top (BR^{-1}B^\top - 2\gamma\sigma\sigma^\top)S_\alpha + 2z_\alpha^\top D^\top S_\alpha + z_\alpha^\top \Gamma^\top Q\Gamma z_\alpha + \text{Tr}(\sigma\sigma^\top \Pi) = 0, \quad (9)$$

where $r_\alpha(T) = z_\alpha^\top(T)\Gamma_f^\top Q_f\Gamma_f z_\alpha(T)$. Under (H2)–(H4), the ODE system (8)–(9) admits a unique solution $(S_\alpha, r_\alpha) \in C([0, T]; \mathbb{R}^n) \times C([0, T]; \mathbb{R})$. Then we obtain the candidate optimal control

$$\bar{u}_\alpha = -R^{-1}B^\top \Pi x_\alpha - R^{-1}B^\top S_\alpha.$$

Now, we show that \bar{u}_α is indeed the optimal control of Problem (BR). Applying Itô's formula to $\langle \Pi x_\alpha, x_\alpha \rangle + 2\langle S_\alpha, x_\alpha \rangle$ and using the completion of squares technique, we have

$$J_\alpha(u_\alpha) = \mathbb{E}\left[e^{\gamma\{\xi_\alpha^\top \Pi(0)\xi_\alpha + 2\xi_\alpha^\top S_\alpha(0) + r_\alpha(0) + \int_0^T \|u_\alpha + R^{-1}B^\top (\Pi x_\alpha + S_\alpha)\|_R^2 dt\}}\right].$$

Since $\gamma > 0$, J_α attains its minimum when $u_\alpha = \bar{u}_\alpha$, which implies that \bar{u}_α is indeed the optimal control. We summarize the above results into the following theorem.

Theorem 3.1. Under (H1)–(H4), the optimal control law of Problem (BR) is given by

$$\bar{u}_\alpha(t) = -R^{-1}(t)B^\top(t)\Pi(t)x_\alpha(t) - R^{-1}(t)B^\top(t)S_\alpha(t), \quad \alpha \in [0, 1], \quad (10)$$

for which the closed-loop state process x_α satisfies

$$dx_\alpha = [(A - BR^{-1}B^\top \Pi)x_\alpha - BR^{-1}B^\top S_\alpha + Dz_\alpha]dt + \sigma dw_\alpha, \quad x_\alpha(0) = \xi_\alpha. \quad (11)$$

Moreover, the optimal cost is given by

$$J_\alpha(\bar{u}_\alpha) = \mathbb{E}\left[\exp\left(\gamma\{\xi_\alpha^\top \Pi(0)\xi_\alpha + 2\xi_\alpha^\top S_\alpha(0) + r_\alpha(0)\}\right)\right]. \quad (12)$$

4 Consistency condition

The best response control problem of Section 3 has been solved by assuming each $z_\alpha(\cdot)$ to be a known function defined on $[0, T]$. Following [10, 21], we specify z_α by imposing a consistency condition such that $z_\alpha(t)$ is regenerated by the graphon weighted average of the individual means:

$$z_\alpha(t) = \int_0^1 g(\alpha, \beta) \mathbb{E}x_\beta(t) d\beta. \quad (13)$$

Combining (11), (8) with (13), we derive the following GMFG equation system:

$$\begin{cases} \dot{z}_\alpha = (A - BR^{-1}B^\top \Pi)z_\alpha + D \int_0^1 g(\alpha, \beta) z_\beta d\beta - BR^{-1}B^\top \int_0^1 g(\alpha, \beta) S_\beta d\beta, \\ \dot{S}_\alpha = -(A^\top - \Pi BR^{-1}B^\top + 2\gamma\Pi\sigma\sigma^\top)S_\alpha + (Q\Gamma - \Pi D)z_\alpha, \\ z_\alpha(0) = \int_0^1 g(\alpha, \beta) m_\beta^x d\beta, \quad S_\alpha(T) = -Q_f\Gamma_f z_\alpha(T), \quad \alpha \in [0, 1], \end{cases} \quad (14)$$

which is a family of fully coupled forward-backward equations. In the remaining part of this section, we employ two methods—the fixed point method and the method of continuity—to investigate the well-posedness of the equation system (14).

4.1 Fixed point method

In this subsection, we use a fixed point method to investigate the solvability of (14). Let $\Psi_z(t, s)$ and $\Psi_S(t, s)$ be the fundamental solution matrices of

$$\dot{y}_1 = (A - BR^{-1}B^\top \Pi)y_1, \quad \text{and} \quad \dot{y}_2 = -(A^\top - \Pi BR^{-1}B^\top + 2\gamma \Pi \sigma \sigma^\top)y_2, \quad (15)$$

with $y_1(t), y_2(t) \in \mathbb{R}^n$ and $\Psi_z(s, s) = \Psi_S(s, s) = I_n$. We transform the solvability problem of (14) to a fixed point problem. We write $z_\alpha(t) = z(\alpha, t)$ as a function of (α, t) and will derive an equation for z_α by eliminating S_α . Let $C([0, 1] \times [0, T]; \mathbb{R}^n)$ be the function space consisting of continuous \mathbb{R}^n -valued functions defined on $[0, 1] \times [0, T]$ with norm $\|x\| = \sup_{\alpha, t} |x(\alpha, t)|$. Define the linear operator Ξ acting on $\tilde{z} \in C([0, 1] \times [0, T]; \mathbb{R}^n)$:

$$\begin{aligned} (\Xi \tilde{z})(\alpha, t) = & \int_0^t \Psi_z(t, s) D \int_0^1 g(\alpha, \beta) \tilde{z}(\beta, s) d\beta ds + \int_0^t \Psi_z(t, s) B R^{-1} B^\top \int_0^1 g(\alpha, \beta) \\ & \times \left\{ \int_s^T \Psi_S(s, r) (Q\Gamma - \Pi D) \tilde{z}(\beta, r) dr - \Psi_S(s, T) Q_f \Gamma_f \tilde{z}(\beta, T) \right\} d\beta ds. \end{aligned}$$

It is straightforward to show that Ξ is from $C([0, 1] \times [0, T]; \mathbb{R}^n)$ to itself and is continuous. Then the solvability of the GMFG equation system (14) reduces to finding a solution $\tilde{z} \in C([0, 1] \times [0, T]; \mathbb{R}^n)$ to the following fixed point equation:

$$\tilde{z}(\alpha, t) = (\Xi \tilde{z})(\alpha, t) + \Psi_z(t, 0) \int_0^1 g(\alpha, \beta) m_\beta^x d\beta.$$

Denote the constants $c_g = \max_{\alpha \in [0, 1]} \int_0^1 g(\alpha, \beta) d\beta$, $c_z = \max_{0 \leq t, s \leq T} |\Psi_z(t, s)|$ and $c_S = \max_{0 \leq t, s \leq T} |\Psi_S(t, s)|$. Let $\|\Xi\|$ denote the norm of Ξ . We obtain

$$\|\Xi\| \leq C_\Xi := c_g c_z |D| T + c_g c_z c_S |B R^{-1} B^\top| \cdot (|Q\Gamma - \Pi D| T + |Q_f \Gamma_f|) T.$$

We have the following solvability result.

Theorem 4.1. Suppose (H1)–(H5) hold. If $C_\Xi < 1$, then the equation system (14) admits a unique solution $(z(\cdot), S(\cdot)) \in C([0, 1] \times [0, T]; \mathbb{R}^n) \times C([0, 1] \times [0, T]; \mathbb{R}^n)$. Furthermore, ODE (9) also admits a unique solution $r(\cdot) \in C([0, 1] \times [0, T]; \mathbb{R})$.

Proof. Under the contraction condition, we obtain a unique solution z from (4.1), which in turns determines a unique solution S_α for (8), and next r_α for (9). \square

Remark 4.2. The contraction condition in the fixed point method may be conservative and typically holds on small time intervals $[0, T]$.

4.2 The method of continuity

This part analyzes solvability by the continuity method under a dominance-monotonicity condition. We need to formulate such a condition for operators acting on an infinite dimensional space. We follow [23] to introduce some notation. Define $L^2[0, 1]$ as the space of all square integrable functions from the vertex set $[0, 1]$ to \mathbb{R} . Let $(L^2[0, 1])^n := L^2[0, 1] \times \cdots \times L^2[0, 1]$ with inner product $\langle \mathbf{a}, \mathbf{b} \rangle := \sum_{i=1}^n \int_0^1 \mathbf{a}_i(\alpha) \mathbf{b}_i(\alpha) d\alpha = \int_0^1 \langle \mathbf{a}(\alpha), \mathbf{b}(\alpha) \rangle d\alpha$, for $\mathbf{a}, \mathbf{b} \in (L^2[0, 1])^n$, where $\mathbf{a}_i \in L^2[0, 1]$ is the i -th component of \mathbf{a} , and $\mathbf{a}(\alpha) = (\mathbf{a}_1(\alpha), \dots, \mathbf{a}_n(\alpha))^\top \in \mathbb{R}^n$ for a fixed index $\alpha \in [0, 1]$. The space $(L^2[0, 1])^n$ with the above inner product is a Hilbert space with norm $\|\mathbf{a}\|_2 = (\int_0^1 |\mathbf{a}(\alpha)|^2 d\alpha)^{\frac{1}{2}}$. Let $\mathcal{L}((L^2[0, 1])^n)$ represent the space of bounded linear operators from $(L^2[0, 1])^n$ to $(L^2[0, 1])^n$ with operator norm $\|\cdot\|_{\text{op}}$. Let $L^2([0, T]; (L^2[0, 1])^n)$ stand for the Hilbert space of equivalent classes of strongly measurable (in the Bochner sense [42, p. 103]) mappings from $[0, T]$ to $(L^2[0, 1])^n$ with

norm $\|\mathbf{d}\|_{L^2([0,T];(L^2[0,1])^n)} = (\int_0^T \|\mathbf{d}(t)\|_2^2 dt)^{\frac{1}{2}}$, and $C([0,T];(L^2[0,1])^n)$ for the Banach space of all continuous mappings from $[0,T]$ to $(L^2[0,1])^n$ with norm

$$\|\mathbf{d}\|_C = \sup_{0 \leq t \leq T} \|\mathbf{d}(t)\|_2. \quad (16)$$

A graphon $\mathbf{G} \in \mathbf{G}_0^{\text{sp}}$ defines a self-adjoint bounded linear operator in $\mathcal{L}(L^2[0,1])$ as follows (we use the notation \mathbf{G} rather than g to emphasize that it is an operator):

$$(\mathbf{G}\mathbf{a}_i)(\alpha) = \int_0^1 \mathbf{G}(\alpha, \beta) \mathbf{a}_i(\beta) d\beta, \quad \forall \alpha \in [0,1], \quad (17)$$

where $\mathbf{G}\mathbf{a}_i \in L^2[0,1]$. Moreover, graphons can be associated with operators from $(L^2[0,1])^n$ to $(L^2[0,1])^n$. We use the square bracket $[\cdot]$ to indicate that the operator is in $\mathcal{L}((L^2[0,1])^n)$. For any $F \in \mathbb{R}^{n \times n}$ and $\mathbf{G} \in \mathbf{G}_0^{\text{sp}}$, the operator $[F\mathbf{G}]$ is defined as $([F\mathbf{G}]\mathbf{a})(\alpha) := F((\mathbf{G}\mathbf{a}_1)(\alpha), \dots, (\mathbf{G}\mathbf{a}_n)(\alpha))^{\top}$, which lies in \mathbb{R}^n for a given α . Since $[F\mathbf{G}] \in \mathcal{L}((L^2[0,1])^n)$, it generates a uniformly continuous (hence strongly continuous) semigroup given by $S_{[F\mathbf{G}]}(t) = \exp(t[F\mathbf{G}]) := \sum_{j=0}^{\infty} \frac{1}{j!} t^j [F\mathbf{G}]^j$, $t \geq 0$ [39, pp. 1-2]. Additionally, $\mathbb{I} \in \mathcal{L}((L^2[0,1])^n)$ denotes the identity operator satisfying $([F\mathbb{I}]\mathbf{a})(\alpha) := F(\mathbf{a}_1(\alpha), \dots, \mathbf{a}_n(\alpha))^{\top} = F\mathbf{a}(\alpha) \in \mathbb{R}^n$.

Now, we rewrite the equation system (14) in the following operator form:

$$\begin{cases} \dot{\mathbf{z}} = (\mathbb{A} + [D\mathbf{G}])\mathbf{z} - [BR^{-1}B^{\top}\mathbf{G}]\mathbf{S}, & \mathbf{z}(0) = \int_0^1 \mathbf{G}(\cdot, \beta) m_{\beta}^x d\beta, \\ \dot{\mathbf{S}} = -(\mathbb{A}^{\top} + [2\gamma\Pi\sigma\sigma^{\top}\mathbb{I}])\mathbf{S} + [(Q\Gamma - \Pi D)\mathbb{I}]\mathbf{z}, & \mathbf{S}(T) = -[Q_f\Gamma_f\mathbb{I}]\mathbf{z}(T), \end{cases} \quad (18)$$

where $\mathbb{A} = [(A - BR^{-1}B^{\top}\Pi)\mathbb{I}]$. In order to investigate the well-posedness of the forward-backward equation system (18), we introduce the dominance-monotonicity condition:

(H6) (Dominance-monotonicity condition). There exist two constants $\mu \geq 0$, $\nu \geq 0$, a matrix $M \in \mathbb{R}^{\tilde{n} \times \tilde{n}}$, and two given matrix value functions $F, H \in L^{\infty}([0,T]; \mathbb{R}^{\tilde{n} \times \tilde{n}})$ (for some $\tilde{n}, \tilde{n} \in \mathbb{N}$) such that the following conditions are fulfilled:

- (i) One of these two cases holds: Case (A) $\mu > 0$ and $\nu = 0$; Case (B) $\mu = 0$ and $\nu > 0$.
- (ii) (Dominance condition). For all $t \in [0, T]$ appearing in $B(t), R(t), D(t), Q(t), \Pi(t), H(t), F(t)$ and all $\mathbf{z}, \mathbf{S} \in (L^2[0,1])^n$,

$$\begin{cases} \|[BR^{-1}B^{\top}\mathbf{G}]\mathbf{S}\|_2 \leq \frac{1}{\mu} \|[H\mathbb{I}]\mathbf{S}\|_2, & \|[Q\Gamma - \Pi D]\mathbb{I}\mathbf{z}\|_2 \leq \frac{1}{\nu} \|[F\mathbb{I}]\mathbf{z}\|_2, \\ \|[Q_f\Gamma_f\mathbb{I}]\mathbf{z}\|_2 \leq \frac{1}{\nu} \|[M\mathbb{I}]\mathbf{z}\|_2. \end{cases}$$

- (iii) (Monotonicity condition). For all $t \in [0, T]$ appearing in $B(t), R(t), D(t), Q(t), \Pi(t), H(t), F(t)$ and all $\mathbf{z}, \mathbf{S} \in (L^2[0,1])^n$,

$$\begin{cases} \langle [D\mathbf{G}]\mathbf{z}, \mathbf{S} \rangle - \langle [BR^{-1}B^{\top}\mathbf{G}]\mathbf{S}, \mathbf{S} \rangle - \langle [2\gamma\Pi\sigma\sigma^{\top}\mathbb{I}]\mathbf{S}, \mathbf{z} \rangle + \langle [(Q\Gamma - \Pi D)\mathbb{I}]\mathbf{z}, \mathbf{z} \rangle \\ \leq -\nu \|[F\mathbb{I}]\mathbf{z}\|_2^2 - \mu \|[H\mathbb{I}]\mathbf{S}\|_2^2, \\ -\langle [Q_f\Gamma_f\mathbb{I}]\mathbf{z}, \mathbf{z} \rangle \geq \nu \|[M\mathbb{I}]\mathbf{z}\|_2^2. \end{cases}$$

We adopt the following convention for condition (ii). When $\mu = 0$ (resp., $\nu = 0$), $\frac{1}{\mu}$ (resp., $\frac{1}{\nu}$) means ∞ . That is, if $\mu = 0$ or $\nu = 0$, then the corresponding dominance condition becomes superfluous and is dropped.

Remark 4.3. In fact, the case with $\mu > 0$ and $\nu > 0$ is covered by both Case (A) and Case (B).

For any $\mathbf{m}, \bar{\mathbf{m}} \in L^2([0, T]; (L^2[0, 1])^n)$ and $\boldsymbol{\eta} \in (L^2[0, 1])^n$, we introduce a family of forward-backward equation systems parameterized by $\kappa \in [0, 1]$:

$$\begin{cases} \dot{\mathbf{z}}^\kappa = \kappa \{ (\mathbb{A} + [D\mathbf{G}])\mathbf{z}^\kappa - [BR^{-1}B^\top \mathbf{G}]\mathbf{S}^\kappa \} - (1 - \kappa)\mu[H^\top H\mathbb{I}]\mathbf{S}^\kappa + \mathbf{m}, \\ \dot{\mathbf{S}}^\kappa = \kappa \{ -(\mathbb{A}^\top + [2\gamma\Pi\sigma\sigma^\top]\mathbb{I})\mathbf{S}^\kappa + [(Q\Gamma - \Pi D)\mathbb{I}]\mathbf{z}^\kappa \} - (1 - \kappa)\nu[F^\top F\mathbb{I}]\mathbf{z}^\kappa + \bar{\mathbf{m}}, \\ \mathbf{z}^\kappa(0) = \int_0^1 \mathbf{G}(\cdot, \beta)m_\beta^x d\beta, \quad \mathbf{S}^\kappa(T) = -\kappa[Q_f\Gamma_f\mathbb{I}]\mathbf{z}^\kappa(T) + (1 - \kappa)\nu[M^\top M\mathbb{I}]\mathbf{z}^\kappa(T) + \boldsymbol{\eta}. \end{cases} \quad (19)$$

We will show that for each $\kappa \in [0, 1]$, the equation system (19) admits a unique solution $(\mathbf{z}^\kappa, \mathbf{S}^\kappa) \in C([0, T]; (L^2[0, 1])^n) \times C([0, T]; (L^2[0, 1])^n)$. It is easy to see that when $\kappa = 1$ and $(\mathbf{m}, \bar{\mathbf{m}}, \boldsymbol{\eta}) = (\mathbf{0}, \mathbf{0}, \mathbf{0})$, equation system (19) reduces to (18). Moreover, when $\kappa = 0$, (19) becomes

$$\begin{cases} \dot{\mathbf{z}}^0 = -\mu[H^\top H\mathbb{I}]\mathbf{S}^0 + \mathbf{m}, \quad \mathbf{z}^0(0) = \int_0^1 \mathbf{G}(\cdot, \beta)m_\beta^x d\beta, \\ \dot{\mathbf{S}}^0 = -\nu[F^\top F\mathbb{I}]\mathbf{z}^0 + \bar{\mathbf{m}}, \quad \mathbf{S}^0(T) = \nu[M^\top M\mathbb{I}]\mathbf{z}^0(T) + \boldsymbol{\eta}. \end{cases} \quad (20)$$

Now, we investigate the existence and uniqueness of a solution to the forward-backward equation (20) in two cases. For Case (A) (i.e., $\mu > 0$ and $\nu = 0$), (20) is in a decoupled form, for which we can solve the backward equation first to get \mathbf{S}^0 , and next obtain \mathbf{z}^0 . For Case (B) (i.e., $\mu = 0$ and $\nu > 0$), (20) is also in a decoupled form and easily solved. Thus, under (H6), (20) admits a unique solution $(\mathbf{z}^0, \mathbf{S}^0) \in C([0, T]; (L^2[0, 1])^n) \times C([0, T]; (L^2[0, 1])^n)$.

The method of continuity works by showing that whenever (19) admits a unique solution for some $\kappa_0 \in [0, 1]$, then unique solvability still holds for all κ slightly large than κ_0 , allowing to cover up to $\kappa = 1$. To develop the continuity arguments, we first give the following a priori estimate of the solution to (19).

Lemma 4.4. Under (H1)–(H6), suppose that $(\mathbf{z}_j^\kappa, \mathbf{S}_j^\kappa) \in C([0, T]; (L^2[0, 1])^n) \times C([0, T]; (L^2[0, 1])^n)$, $j = 1, 2$, are solutions of (19) corresponding to $(\mathbf{m}_j, \bar{\mathbf{m}}_j, \boldsymbol{\eta}_j)$, respectively. Then one has the following estimate

$$\|\delta\mathbf{z}^\kappa\|_C^2 + \|\delta\mathbf{S}^\kappa\|_C^2 \leq K \left[\|\delta\boldsymbol{\eta}\|_2^2 + \int_0^T \|\delta\mathbf{m}(t)\|_2^2 dt + \int_0^T \|\delta\bar{\mathbf{m}}(t)\|_2^2 dt \right],$$

where the norm $\|\cdot\|_C$ is defined in (16). Here, $\delta\mathbf{z}^\kappa = \mathbf{z}_1^\kappa - \mathbf{z}_2^\kappa$ and other terms are similarly defined. K is a positive constant depending on T , μ , ν and the upper bounds of all coefficients in (1) and (3), Π , the norm of H , F and M , and $\|\mathbf{G}\|_{\text{op}}$.

Proof. We present the proofs for Case (A) and Case (B) separately. For Case (A) (i.e., $\mu > 0$ and $\nu = 0$), according to (19) and (H6)–(ii), it holds that

$$\begin{aligned} \|\delta\mathbf{z}^\kappa\|_C^2 &\leq K_1 \left(\int_0^T \|\delta\mathbf{m}(t)\|_2^2 dt + \int_0^T \|[H\mathbb{I}]\delta\mathbf{S}^\kappa(t)\|_2^2 dt \right), \\ \|\delta\mathbf{S}^\kappa\|_C^2 &\leq K_1 \left(\int_0^T \|\delta\bar{\mathbf{m}}(t)\|_2^2 dt + \|\delta\mathbf{z}^\kappa\|_C^2 + \|\delta\boldsymbol{\eta}\|_2^2 \right), \end{aligned}$$

where $K_1 > 0$ is a constant. Eliminating $\|\delta\mathbf{z}^\kappa\|_C^2$ in the second inequality, we obtain

$$\|\delta\mathbf{z}^\kappa\|_C^2 + \|\delta\mathbf{S}^\kappa\|_C^2 \leq K_2 \left[\|\delta\boldsymbol{\eta}\|_2^2 + \int_0^T \left\{ \|\delta\mathbf{m}(t)\|_2^2 + \|\delta\bar{\mathbf{m}}(t)\|_2^2 + \|[H\mathbb{I}]\delta\mathbf{S}^\kappa(t)\|_2^2 \right\} dt \right], \quad (21)$$

where $K_2 > 0$ is a constant. Recalling (19), we derive the equation of $(\delta\mathbf{z}^\kappa, \delta\mathbf{S}^\kappa)$. Taking the differential of $\langle \delta\mathbf{z}^\kappa, \delta\mathbf{S}^\kappa \rangle$ and integrating both sides over $[0, T]$, we then use the monotonicity condition (H6)–(iii)

to obtain

$$\begin{aligned}
& \langle \delta \mathbf{z}^\kappa(T), \delta \mathbf{S}^\kappa(T) \rangle - \langle \delta \mathbf{z}^\kappa(0), \delta \mathbf{S}^\kappa(0) \rangle \\
&= \int_0^T \left\{ \langle \delta \mathbf{z}^\kappa, \kappa [-(\mathbb{A}^\top + [2\gamma \Pi \sigma \sigma^\top \mathbb{I}]) \delta \mathbf{S}^\kappa + [(Q\Gamma - D^\top \Pi) \mathbb{I}] \delta \mathbf{z}^\kappa] + \delta \bar{\mathbf{m}} \rangle \right. \\
&\quad \left. + \langle \kappa [(\mathbb{A} + [D\mathbf{G}]) \delta \mathbf{z}^\kappa - [BR^{-1}B^\top \mathbf{G}] \delta \mathbf{S}^\kappa] - (1 - \kappa) \mu [H^\top H \mathbb{I}] \delta \mathbf{S}^\kappa + \delta \mathbf{m}, \delta \mathbf{S}^\kappa \rangle \right\} dt \\
&\leq \int_0^T \left(-\mu \|[H\mathbb{I}] \delta \mathbf{S}^\kappa(t)\|_2^2 + \langle \delta \mathbf{m}(t), \delta \mathbf{S}^\kappa(t) \rangle + \langle \delta \mathbf{z}^\kappa(t), \delta \bar{\mathbf{m}}(t) \rangle \right) dt,
\end{aligned} \tag{22}$$

and

$$\langle \delta \mathbf{z}^\kappa(T), \delta \mathbf{S}^\kappa(T) \rangle = \langle \delta \mathbf{z}^\kappa(T), -\kappa [Q_f \Gamma_f \mathbb{I}] \delta \mathbf{z}^\kappa(T) + \delta \boldsymbol{\eta} \rangle \geq \langle \delta \mathbf{z}^\kappa(T), \delta \boldsymbol{\eta} \rangle. \tag{23}$$

Then we have

$$\mu \int_0^T \|[H\mathbb{I}] \delta \mathbf{S}^\kappa(t)\|_2^2 dt \leq \int_0^T \left(\langle \delta \mathbf{m}(t), \delta \mathbf{S}^\kappa(t) \rangle + \langle \delta \mathbf{z}^\kappa(t), \delta \bar{\mathbf{m}}(t) \rangle \right) dt - \langle \delta \mathbf{z}^\kappa(T), \delta \boldsymbol{\eta} \rangle,$$

which implies

$$\begin{aligned}
& \int_0^T \|[H\mathbb{I}] \delta \mathbf{S}^\kappa(t)\|_2^2 dt \\
&\leq \frac{1}{\mu} \int_0^T \left(\langle \delta \mathbf{m}(t), \delta \mathbf{S}^\kappa(t) \rangle + \langle \delta \mathbf{z}^\kappa(t), \delta \bar{\mathbf{m}}(t) \rangle \right) dt - \frac{1}{\mu} \langle \delta \mathbf{z}^\kappa(T), \delta \boldsymbol{\eta} \rangle \\
&\leq \frac{1}{\mu} \left(\|\delta \mathbf{S}^\kappa\|_C \cdot \int_0^T \|\delta \mathbf{m}(t)\|_2 dt + \|\delta \mathbf{z}^\kappa\|_C \cdot \int_0^T \|\delta \bar{\mathbf{m}}(t)\|_2 dt + \|\delta \mathbf{z}^\kappa\|_C \cdot \|\delta \boldsymbol{\eta}\|_2 \right) \\
&\leq \frac{1}{4\mu\theta_1} \left(\|\delta \boldsymbol{\eta}\|_2^2 + T \int_0^T \left\{ \|\delta \mathbf{m}(t)\|_2^2 + \|\delta \bar{\mathbf{m}}(t)\|_2^2 \right\} dt \right) + \frac{2\theta_1}{\mu} \left(\|\delta \mathbf{S}^\kappa\|_C^2 + \|\delta \mathbf{z}^\kappa\|_C^2 \right),
\end{aligned} \tag{24}$$

where $\theta_1 > 0$ is arbitrary. We now select $\theta_1 = \frac{\mu}{4K_2}$, where K_2 is from (21). Combining (21) with (24), we obtain the desired result.

For Case (B) (i.e., $\mu = 0$ and $\nu > 0$), according to (19) and (H6)–(ii), we have

$$\begin{aligned}
\|\delta \mathbf{z}^\kappa\|_C^2 &\leq K_3 \left(\int_0^T \|\delta \mathbf{m}(t)\|_2^2 dt + \|\delta \mathbf{S}^\kappa\|_C^2 \right), \\
\|\delta \mathbf{S}^\kappa\|_C^2 &\leq K_3 \left(\int_0^T \|\delta \bar{\mathbf{m}}(t)\|_2^2 dt + \int_0^T \|[F\mathbb{I}] \delta \mathbf{z}^\kappa(t)\|_2^2 dt + \|[M\mathbb{I}] \delta \mathbf{z}^\kappa(T)\|_2^2 + \|\delta \boldsymbol{\eta}\|_2^2 \right),
\end{aligned}$$

where $K_3 > 0$ is a constant. Eliminating $\|\delta \mathbf{S}^\kappa\|_C^2$ in the first inequality, then

$$\begin{aligned}
& \|\delta \mathbf{z}^\kappa\|_C^2 + \|\delta \mathbf{S}^\kappa\|_C^2 \\
&\leq K_4 \left[\|\delta \boldsymbol{\eta}\|_2^2 + \int_0^T \left\{ \|\delta \mathbf{m}(t)\|_2^2 + \|\delta \bar{\mathbf{m}}(t)\|_2^2 + \|[F\mathbb{I}] \delta \mathbf{z}^\kappa(t)\|_2^2 \right\} dt + \|[M\mathbb{I}] \delta \mathbf{z}^\kappa(T)\|_2^2 \right],
\end{aligned} \tag{25}$$

where $K_4 > 0$ is a constant. Similar to (22)–(23), and using the monotonicity condition (H6)–(iii), we have

$$\begin{aligned}
& \langle \delta \mathbf{z}^\kappa(T), \delta \mathbf{S}^\kappa(T) \rangle - \langle \delta \mathbf{z}^\kappa(0), \delta \mathbf{S}^\kappa(0) \rangle \\
&= \int_0^T \left\{ \langle \delta \mathbf{z}^\kappa, \kappa [-(\mathbb{A}^\top + [2\gamma \Pi \sigma \sigma^\top \mathbb{I}]) \delta \mathbf{S}^\kappa + [(Q\Gamma - D^\top \Pi) \mathbb{I}] \delta \mathbf{z}^\kappa] - (1 - \kappa) \nu [F^\top F \mathbb{I}] \delta \mathbf{z}^\kappa + \delta \bar{\mathbf{m}} \rangle \right. \\
&\quad \left. + \langle \kappa [(\mathbb{A} + [D\mathbf{G}]) \delta \mathbf{z}^\kappa - [BR^{-1}B^\top \mathbf{G}] \delta \mathbf{S}^\kappa] + \delta \mathbf{m}, \delta \mathbf{S}^\kappa \rangle \right\} dt \\
&\leq \int_0^T \left(-\nu \|[F\mathbb{I}] \delta \mathbf{z}^\kappa(t)\|_2^2 + \langle \delta \mathbf{m}(t), \delta \mathbf{S}^\kappa(t) \rangle + \langle \delta \mathbf{z}^\kappa(t), \delta \bar{\mathbf{m}}(t) \rangle \right) dt,
\end{aligned}$$

and

$$\begin{aligned}\langle \delta \mathbf{z}^\kappa(T), \delta \mathbf{S}^\kappa(T) \rangle &= \langle \delta \mathbf{z}^\kappa(T), -\kappa[Q_f \Gamma_f \mathbb{I}] \delta \mathbf{z}^\kappa(T) + (1 - \kappa)\nu[M^\top M \mathbb{I}] \delta \mathbf{z}^\kappa(T) + \delta \boldsymbol{\eta} \rangle \\ &\geq \nu \|[M \mathbb{I}] \delta \mathbf{z}^\kappa(T)\|_2^2 + \langle \delta \mathbf{z}^\kappa(T), \delta \boldsymbol{\eta} \rangle.\end{aligned}$$

Then we have

$$\begin{aligned}&\nu \|[M \mathbb{I}] \delta \mathbf{z}^\kappa(T)\|_2^2 + \nu \int_0^T \|[F \mathbb{I}] \delta \mathbf{z}^\kappa(t)\|_2^2 dt \\ &\leq \int_0^T \left(\langle \delta \mathbf{m}(t), \delta \mathbf{S}^\kappa(t) \rangle + \langle \delta \mathbf{z}^\kappa(t), \delta \bar{\mathbf{m}}(t) \rangle \right) dt - \langle \delta \mathbf{z}^\kappa(T), \delta \boldsymbol{\eta} \rangle,\end{aligned}$$

which implies

$$\begin{aligned}&\|[M \mathbb{I}] \delta \mathbf{z}^\kappa(T)\|_2^2 + \int_0^T \|[F \mathbb{I}] \delta \mathbf{z}^\kappa(t)\|_2^2 dt \\ &\leq \frac{1}{\nu} \int_0^T \left(\langle \delta \mathbf{m}(t), \delta \mathbf{S}^\kappa(t) \rangle + \langle \delta \mathbf{z}^\kappa(t), \delta \bar{\mathbf{m}}(t) \rangle \right) dt - \frac{1}{\nu} \langle \delta \mathbf{z}^\kappa(T), \delta \boldsymbol{\eta} \rangle \\ &\leq \frac{1}{\nu} \left(\|\delta \mathbf{S}^\kappa\|_C \cdot \int_0^T \|\delta \mathbf{m}(t)\|_2 dt + \|\delta \mathbf{z}^\kappa\|_C \cdot \int_0^T \|\delta \bar{\mathbf{m}}(t)\|_2 dt + \|\delta \mathbf{z}^\kappa\|_C \cdot \|\delta \boldsymbol{\eta}\|_2 \right) \\ &\leq \frac{1}{4\nu\theta_2} \left(\|\delta \boldsymbol{\eta}\|_2^2 + T \int_0^T \left\{ \|\delta \mathbf{m}(t)\|_2^2 + \|\delta \bar{\mathbf{m}}(t)\|_2^2 \right\} dt \right) + \frac{2\theta_2}{\nu} \left(\|\delta \mathbf{S}^\kappa\|_C^2 + \|\delta \mathbf{z}^\kappa\|_C^2 \right),\end{aligned}\tag{26}$$

where $\theta_2 > 0$ is arbitrary. We now select $\theta_2 = \frac{\mu}{4K_4}$, where K_4 is from (25). Combining (25) with (26), we obtain the desired result. \square

With the aid of Lemma 4.4, we obtain the following lemma.

Lemma 4.5. Under (H1)–(H6), there exists a constant $\delta_0 > 0$ such that if the equation system (19) is uniquely solvable for some $\kappa_0 \in [0, 1]$ whenever $\boldsymbol{\eta} \in (L^2[0, 1])^n$ and $\mathbf{m}, \bar{\mathbf{m}} \in L^2([0, T]; (L^2[0, 1])^n)$, then the same conclusion holds for all $\kappa \in [\kappa_0, (\kappa_0 + \delta_0) \wedge 1]$, where δ_0 does not depend on κ_0 .

Proof. Take any $\bar{\delta} \in (0, \delta_0]$, where $\delta_0 > 0$ will be selected later. For any $(\mathbf{z}, \mathbf{S}) \in C([0, T]; (L^2[0, 1])^n) \times C([0, T]; (L^2[0, 1])^n)$, consider the following forward-backward equation system

$$\begin{cases} \dot{\tilde{\mathbf{z}}} = \kappa_0 [(\mathbb{A} + [D\mathbf{G}])\tilde{\mathbf{z}} - [BR^{-1}B^\top \mathbf{G}]\tilde{\mathbf{S}}] - (1 - \kappa_0)\mu[H^\top H \mathbb{I}]\tilde{\mathbf{S}} + \mathbf{n}, \\ \dot{\tilde{\mathbf{S}}} = \kappa_0 [-(\mathbb{A}^\top + [2\gamma\Pi\sigma\sigma^\top \mathbb{I}])\tilde{\mathbf{S}} + [(Q\Gamma - \Pi D)\mathbb{I}]\tilde{\mathbf{z}}] - (1 - \kappa_0)\nu[F^\top F \mathbb{I}]\tilde{\mathbf{z}} + \bar{\mathbf{n}}, \\ \tilde{\mathbf{z}}(0) = \int_0^1 \mathbf{G}(\cdot, \beta) m_\beta^x d\beta, \quad \tilde{\mathbf{S}}(T) = -\kappa_0[Q_f \Gamma_f \mathbb{I}]\tilde{\mathbf{z}}(T) + (1 - \kappa_0)\nu[M^\top M \mathbb{I}]\tilde{\mathbf{z}}(T) + \bar{\boldsymbol{\eta}}, \end{cases}\tag{27}$$

where

$$\begin{aligned}\mathbf{n} &= \mathbf{m} + \bar{\delta} [(\mathbb{A} + [D\mathbf{G}])\mathbf{z} - [BR^{-1}B^\top \mathbf{G}]\mathbf{S}] + \bar{\delta}\mu[H^\top H \mathbb{I}]\mathbf{S}, \\ \bar{\mathbf{n}} &= \bar{\mathbf{m}} + \bar{\delta} [-(\mathbb{A}^\top + [2\gamma\Pi\sigma\sigma^\top \mathbb{I}])\mathbf{S} + [(Q\Gamma - \Pi D)\mathbb{I}]\mathbf{z}] + \bar{\delta}\nu[F^\top F \mathbb{I}]\mathbf{S}, \\ \bar{\boldsymbol{\eta}} &= \boldsymbol{\eta} - \bar{\delta}[Q_f \Gamma_f \mathbb{I}]\mathbf{z}(T) - \bar{\delta}\nu[M^\top M \mathbb{I}]\mathbf{z}(T).\end{aligned}$$

Here $\mathbf{m}, \bar{\mathbf{m}}$ and $\boldsymbol{\eta}$ take the same form as in (19). We can easily check that $\bar{\boldsymbol{\eta}} \in (L^2[0, 1])^n$ and $\mathbf{n}, \bar{\mathbf{n}} \in L^2([0, T]; (L^2[0, 1])^n)$. Then by our assumption, (27) admits a unique solution $(\tilde{\mathbf{z}}, \tilde{\mathbf{S}}) \in C([0, T]; (L^2[0, 1])^n) \times C([0, T]; (L^2[0, 1])^n)$ for some $\kappa_0 \in [0, 1]$ whenever $\boldsymbol{\eta} \in (L^2[0, 1])^n$ and $\mathbf{m}, \bar{\mathbf{m}} \in L^2([0, T]; (L^2[0, 1])^n)$. Thus we have a well-defined mapping from (\mathbf{z}, \mathbf{S}) to $(\tilde{\mathbf{z}}, \tilde{\mathbf{S}})$, which is denoted by

$$(\tilde{\mathbf{z}}, \tilde{\mathbf{S}}) = \mathbf{I}_{\kappa_0 + \bar{\delta}}(\mathbf{z}, \mathbf{S}).$$

Next, we will show that, if $\bar{\delta}$ is sufficiently small, the operator $\mathbf{I}_{\kappa_0 + \bar{\delta}}$ from $C([0, T]; (L^2[0, 1])^n) \times C([0, T]; (L^2[0, 1])^n)$ to itself is a contraction. Let $(\tilde{\mathbf{z}}_1, \tilde{\mathbf{S}}_1)$ be another solution of (27), when (\mathbf{z}, \mathbf{S}) is replaced by $(\mathbf{z}_1, \mathbf{S}_1)$. Denote $\delta \tilde{\mathbf{z}} = \tilde{\mathbf{z}}_1 - \tilde{\mathbf{z}}$ and $\delta \tilde{\mathbf{S}} = \tilde{\mathbf{S}}_1 - \tilde{\mathbf{S}}$. Following the proving method of

Lemma 4.4, we can show $\|(\delta\tilde{\mathbf{z}}, \delta\tilde{\mathbf{S}})\|_C^2 \leq K_5 \bar{\delta}^2 \|(\delta\mathbf{z}, \delta\mathbf{S})\|_C^2$, where K_5 is a positive constant independent of κ_0 and $\bar{\delta}$. Thus we can choose $\delta_0 = \frac{1}{2\sqrt{K_5}}$, so that for any $\bar{\delta} \in (0, \delta_0]$ satisfying $\kappa_0 + \bar{\delta} \leq 1$, we have $\|(\delta\tilde{\mathbf{z}}, \delta\tilde{\mathbf{S}})\|_C^2 \leq \frac{1}{2} \|(\delta\mathbf{z}, \delta\mathbf{S})\|_C^2$, which implies that the operator $\mathbf{I}_{\kappa_0 + \bar{\delta}}$ is a contraction. Therefore, it has a unique fixed point, which is the unique solution of (19) with $\kappa = \kappa_0 + \bar{\delta}$. \square

We have the following existence result.

Theorem 4.6. Under (H1)–(H6), the forward-backward equation system (18) admits a unique solution $(\mathbf{z}, \mathbf{S}) \in C([0, T]; (L^2[0, 1])^n) \times C([0, T]; (L^2[0, 1])^n)$.

Proof. From (20), we know that (19) admits a unique solution when $\kappa = \kappa_0 = 0$. It then follows from Lemma 4.5 that there exists a positive constant δ_0 such that for each $\kappa \in [\kappa_0, (\kappa_0 + \delta_0) \wedge 1]$, (19) admits a unique solution. Thus, we can repeat this process ι times with $1 \leq \iota\delta_0 < 1 + \delta_0$. It then follows that, in particular, the forward-backward equation system (19) admits a unique solution when $\kappa = 1$ and $(\mathbf{m}, \bar{\mathbf{m}}, \boldsymbol{\eta}) = (\mathbf{0}, \mathbf{0}, \mathbf{0})$. \square

4.3 Example

For illustration of the monotonicity condition (H6)–(iii), we give the following example.

Example 4.7. Each graphon \mathbf{G} gives a self-adjoint compact operator from $L^2[0, 1]$ to $L^2[0, 1]$ [34, Section 7.5], and hence has the following discrete spectral decomposition

$$\mathbf{G}(\alpha, \beta) = \sum_{\ell \in \mathcal{I}_\lambda} \lambda_\ell \mathbf{f}_\ell(\alpha) \mathbf{f}_\ell(\beta), \quad (28)$$

where $\mathbf{f}_\ell(\cdot) \in L^2[0, 1]$, $\{\lambda_\ell\}_{\ell \geq 1} \subset \mathbb{R}$ is the set of eigenvalues, and the convergence is in the $(L^2[0, 1])^n$ sense. Here, $\{\mathbf{f}_\ell\}_{\ell \in \mathcal{I}_\lambda}$ represents the corresponding set of orthonormal eigenfunctions, and \mathcal{I}_λ is the index multiset for all the nonzero eigenvalues of \mathbf{G} . Since the graphon operator \mathbf{G} defined in (17) is a Hilbert-Schmidt integral operator and hence a compact operator in $\mathcal{L}(L^2[0, 1])$, the number of elements in \mathcal{I}_λ is either finite or countably infinite; see [34, Section 7.5]. In particular, if $\{\lambda_1, \lambda_2, \dots\}$ is a countable multiset of nonzero eigenvalues, then $\lim_{\ell \rightarrow \infty} \lambda_\ell = 0$, and every nonzero eigenvalue has finite multiplicity.

Now we consider a graphon with spectral decomposition (28) and denote the index set of all strictly positive eigenvalues λ_ℓ by \mathcal{I}_λ^+ . We further assume that \mathcal{I}_λ^+ is finite. Without loss of generality, we assume $\mathcal{I}_\lambda^+ = \{\lambda_1, \dots, \lambda_d\}$ and $\lambda_{\min} = \min_{1 \leq \ell \leq d} \lambda_\ell$. Let \mathcal{H} be the subspace generated by the orthonormal basis functions $\{\mathbf{f}_1, \dots, \mathbf{f}_d\}$ and \mathcal{H}^\perp be the complement subspace in $L^2[0, 1]$. Assume

$$m^x \in \mathcal{H}^n := \mathcal{H} \times \dots \times \mathcal{H}. \quad (29)$$

In view of the proof of Lemma 4.5, the existence and uniqueness analysis can be carried out with the solution lying in \mathcal{H}^n subject to (29). Then we only need to establish the monotonicity condition for $\mathbf{z}, \mathbf{S} \in \mathcal{H}^n$. Moreover, we assume that for every $t \in [0, T]$, the following inequalities hold:

$$\begin{aligned} & -\frac{Q\Gamma + \Gamma^\top Q - \Pi D - D^\top \Pi}{2} - \gamma^2 \Pi \sigma \sigma^\top \sigma^\top \sigma \Pi - \frac{DD^\top}{4} \geq 0, \\ & BR^{-1}B^\top \lambda_{\min} - 2I_n \gg 0, \quad -(Q_f \Gamma_f + \Gamma_f^\top Q_f) \geq 0, \end{aligned} \quad (30)$$

where Π is determined by (7). Let $\nu = \inf_{0 \leq t \leq T} |\tilde{\lambda}_Q^*(t) \vee \tilde{\lambda}_{Q_f}^*|$ and $\mu = \inf_{0 \leq t \leq T} |\tilde{\lambda}_R^*(t)|$, where $\tilde{\lambda}_Q^*(t)$ (resp., $\tilde{\lambda}_R^*(t)$, $\tilde{\lambda}_{Q_f}^*(t)$) is the largest eigenvalue of the symmetric matrix $\frac{Q(t)\Gamma + \Gamma^\top Q(t) - D^\top(t)\Pi(t) - \Pi(t)D(t)}{2} + \gamma^2 \Pi(t)\sigma(t)\sigma^\top(t)\sigma^\top(t)\sigma(t)\Pi(t) + \frac{D(t)D^\top(t)}{4}$ (resp., $2I_n - B(t)R^{-1}(t)B^\top(t)\lambda_{\min}$, $\frac{Q_f \Gamma_f + \Gamma_f^\top Q_f}{2}$). By (30),

we obtain $\mu > 0$, then

$$\begin{aligned}
& \langle [D\mathbf{G}]\mathbf{z}, \mathbf{S} \rangle - \langle [BR^{-1}B^\top \mathbf{G}]\mathbf{S}, \mathbf{S} \rangle - \langle [2\gamma\Pi\sigma\sigma^\top \mathbb{I}]\mathbf{S}, \mathbf{z} \rangle + \langle [(Q\Gamma - \Pi D)\mathbb{I}]\mathbf{z}, \mathbf{z} \rangle \\
&= \langle [(Q\Gamma - \Pi D)\mathbb{I}]\mathbf{z}, \mathbf{z} \rangle - \langle [BR^{-1}B^\top \mathbf{G}]\mathbf{S}, \mathbf{S} \rangle \\
&\quad - 2\gamma\Pi\sigma\sigma^\top \int_0^1 \mathbf{S}(\alpha)\mathbf{z}(\alpha)d\alpha + D \int_0^1 \int_0^1 \mathbf{G}(\alpha, \beta)\mathbf{z}(\beta)d\beta\mathbf{S}(\alpha)d\alpha \\
&\leq \left\langle \left[\left(Q\Gamma - \Pi D + \gamma^2\Pi\sigma\sigma^\top \sigma^\top \sigma\Pi + \frac{DD^\top}{4} \right) \mathbb{I} \right] \mathbf{z}, \mathbf{z} \right\rangle + \langle [(2I_n - BR^{-1}B^\top \lambda_{\min})\mathbb{I}]\mathbf{S}, \mathbf{S} \rangle \\
&\leq \tilde{\lambda}_Q^*(t)\|\mathbf{z}\|_2^2 + \tilde{\lambda}_R^*(t)\|\mathbf{S}\|_2^2 \\
&\leq -\nu\|\mathbf{z}\|_2^2 - \mu\|\mathbf{S}\|_2^2,
\end{aligned}$$

for all $\mathbf{z}, \mathbf{S} \in \mathcal{H}^n$, and moreover,

$$-\langle [Q_f\Gamma_f\mathbb{I}]\mathbf{z}, \mathbf{z} \rangle \geq -\tilde{\lambda}_{Q_f}^*\|\mathbf{z}\|_2^2 \geq \nu\|\mathbf{z}\|_2^2.$$

Thus the desired monotonicity condition is verified.

Remark 4.8. i) Example 4.7 shows how to construct concrete models to verify the monotonicity condition. In fact, even if $d := |\mathcal{I}_\lambda^+| = \infty$, as long as m^x remains in a fixed finite dimensional subspace of the space spanned by $\{\mathbf{f}_\ell, \ell \in \mathcal{I}_\lambda^+\}$, the monotonicity condition can be verified provided that inequalities (30) hold.

ii) In the risk-neutral case (i.e., the limiting case of (14) with $\gamma \rightarrow 0$), the constraints on the initial mean m^x imposed in Example 4.7 are no longer required. For example, we consider the uniform attachment graphs, which converge to the limit graphon $\mathbf{G}(\alpha, \beta) = 1 - \max(\alpha, \beta)$, $\alpha, \beta \in [0, 1]$, under the cut metric with probability one (see [34, Proposition 11.40]). The spectral decomposition of \mathbf{G} is given by

$$\mathbf{G}(\alpha, \beta) = \sum_{k=1,3,5,\dots} \frac{4}{k^2\pi^2} \sqrt{2} \cos\left(\frac{\pi k\alpha}{2}\right) \times \sqrt{2} \cos\left(\frac{\pi k\beta}{2}\right).$$

Here, the eigenvalues of \mathbf{G} are given by $\lambda_k = \frac{4}{k^2\pi^2}$ for positive odd integers k , and $\lambda_k = 0$ for positive even integers k . Then for every $t \in [0, T]$, if $D = 0$, $-(Q\Gamma + \Gamma^\top Q) \geq 0$, $-(Q_f\Gamma_f + \Gamma_f^\top Q_f) \geq 0$, we can easily check that equation system (18) satisfies Case (A) for the monotonicity condition. To show this, let λ_{\min} be the smallest eigenvalue of graphon \mathbf{G} (here $\lambda_{\min} = 0$), $\nu = \inf_{0 \leq t \leq T} |\lambda_Q^*(t) \vee \lambda_{Q_f}^*|$, $\mu = \inf_{0 \leq t \leq T} |\lambda_R^*(t)|$, where $\lambda_Q^*(t)$ (resp., $\lambda_R^*(t)$, $\lambda_{Q_f}^*(t)$) is the largest eigenvalue of the symmetric matrix $\frac{Q(t)\Gamma + \Gamma^\top Q(t)}{2}$ (resp., $-R^{-1}(t)$, $\frac{Q_f\Gamma_f + \Gamma_f^\top Q_f}{2}$). Subsequently, we can verify $\mu > 0$,

$$-\langle [Q_f\Gamma_f\mathbb{I}]\mathbf{z}, \mathbf{z} \rangle \geq -\lambda_{Q_f}^*\|\mathbf{z}\|_2^2 \geq \nu\|\mathbf{z}\|_2^2$$

and

$$\begin{aligned}
& \langle [D\mathbf{G}]\mathbf{z}, \mathbf{S} \rangle - \langle [BR^{-1}B^\top \mathbf{G}]\mathbf{S}, \mathbf{S} \rangle - \langle [2\gamma\Pi\sigma\sigma^\top \mathbb{I}]\mathbf{S}, \mathbf{z} \rangle + \langle [(Q\Gamma - \Pi D)\mathbb{I}]\mathbf{z}, \mathbf{z} \rangle \\
&\leq \langle [Q\Gamma\mathbb{I}]\mathbf{z}, \mathbf{z} \rangle - \langle R^{-1}[\sqrt{\lambda_{\min}}B\mathbb{I}]\mathbf{S}, [\sqrt{\lambda_{\min}}B\mathbb{I}]\mathbf{S} \rangle \\
&\leq \lambda_Q^*(t)\|\mathbf{z}\|_2^2 + \lambda_R^*(t)\|[\sqrt{\lambda_{\min}}B\mathbb{I}]\mathbf{S}\|_2^2 \\
&\leq -\nu\|\mathbf{z}\|_2^2 - \mu\|[\sqrt{\lambda_{\min}}B\mathbb{I}]\mathbf{S}\|_2^2.
\end{aligned}$$

Hence the monotonicity condition is verified.

5 ε -Nash equilibrium

Suppose that (z, S) has been determined by Theorem 4.6. Taking the resulting function $S_\alpha(t)$ for the best response control law (10), we construct the decentralized individual strategies in the N -player

model by appropriately matching the parameter α . Specifically, we construct the set of decentralized strategies $\hat{u} = (\hat{u}_1, \dots, \hat{u}_N)$ with

$$\hat{u}_i = -R^{-1}B^\top \Pi \hat{x}_i - R^{-1}B^\top S_{I_i^*}, \quad (31)$$

where I_i^* is the midpoint of the subinterval $I_i \in \{I_1, \dots, I_N\}$ of length $\frac{1}{N}$ (see Section 2.1), as the nodal index of agent \mathcal{A}_i in the N -player model. The resulting closed-loop dynamics for \hat{x}_i are governed by

$$d\hat{x}_i = [(A - BR^{-1}B^\top \Pi)\hat{x}_i + D\hat{x}_i^{(N)} - BR^{-1}B^\top S_{I_i^*}]dt + \sigma dw_i, \quad \hat{x}_i(0) = \xi_i, \quad (32)$$

where $\hat{x}_i^{(N)} = \frac{1}{N} \sum_{j=1}^N g_{ij}^N \hat{x}_j$.

To analyze the performance of \hat{u} in terms of an ε -Nash equilibrium, we introduce the following definition.

Definition 5.1. The set of strategies $\hat{u} = (\hat{u}_1, \dots, \hat{u}_N)$ is called an ε -Nash equilibrium with respect to costs \mathcal{J}_i , $1 \leq i \leq N$, if for $\varepsilon \geq 0$, the N inequalities hold:

$$\mathcal{J}_i(\hat{u}_i, \hat{u}_{-i}) \leq \inf_{u_i \in \mathcal{U}_c^i} \mathcal{J}_i(u_i, \hat{u}_{-i}) + \varepsilon, \quad \text{for all } 1 \leq i \leq N,$$

where \mathcal{U}_c^i is defined by (2) as the centralized strategy set for agent \mathcal{A}_i .

We need to introduce the following assumption, which specifies the nature of the approximation error between g^N for the finite graph and the graphon function g .

(H7) The sequence $\{g^N, N \geq 2\}$ and the graphon limit g satisfy

$$\lim_{N \rightarrow \infty} \max_{1 \leq i \leq N} \sum_{j=1}^N \left| \frac{g_{ij}^N}{N} - \int_{\beta \in I_j} g(I_i^*, \beta) d\beta \right| = 0.$$

For further estimate of (32), we need to introduce an auxiliary process. By taking the nodal parameter in (10) as $\alpha = I_i^* \in [0, 1]$, we denote the best response control law

$$\bar{u}_i = -R^{-1}B^\top \Pi \bar{x}_i - R^{-1}B^\top S_{I_i^*}, \quad (33)$$

and accordingly introduce the mean-field limit dynamics

$$d\bar{x}_i = [(A - BR^{-1}B^\top \Pi)\bar{x}_i + D\bar{z}_i - BR^{-1}B^\top S_{I_i^*}]dt + \sigma dw_i, \quad \bar{x}_i(0) = \xi_i. \quad (34)$$

Here, $\bar{z}_i = z_{I_i^*}$ is intended to approximate the weighted state-average $\hat{x}_i^{(N)}$ at node i .

Let us introduce the following notation:

$$\begin{aligned} \varepsilon_{1,N} &= \max_{1 \leq i \leq N} \sum_{j=1}^N \left| \frac{g_{ij}^N}{N} - \int_{\beta \in I_j} g(I_i^*, \beta) d\beta \right|, \\ \varepsilon_{2,N} &= \sup_{0 \leq \alpha \leq 1} \sup_{0 \leq t \leq T} \left| z_\alpha^N(t) - z_\alpha(t) \right|, \\ \varepsilon_{3,N} &= \sup_{0 \leq \alpha \leq 1} \left| \mathbb{E} \xi_\alpha^N - m_\alpha^x \right|, \end{aligned}$$

where $z_\alpha^N = \sum_{i=1}^N \mathbb{1}_{I_i}(\alpha) z_i$ and $\xi_\alpha^N = \sum_{i=1}^N \mathbb{1}_{I_i}(\alpha) \xi_i$. Based on (H1), (H7) and the uniform continuity of $z_\alpha(t)$ with respect to $\alpha \in [0, 1]$, it follows that $\lim_{N \rightarrow \infty} (\varepsilon_{1,N} + \varepsilon_{2,N} + \varepsilon_{3,N}) = 0$. Then we have the following estimates.

Lemma 5.2. Under (H1)–(H7), there exists a constant C_0 independent of N such that

$$\sup_{1 \leq i \leq N} \sup_{0 \leq t \leq T} |\hat{x}_i^{(N)} - z_i| \quad (35)$$

$$\leq \frac{C_0}{N} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right| + \frac{C_0}{N} \sup_{0 \leq t \leq T} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right| + C_0 \varepsilon_N,$$

$$\sup_{1 \leq i \leq N} \sup_{0 \leq t \leq T} |\hat{x}_i - \bar{x}_i| \quad (36)$$

$$\leq \frac{C_0}{N} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right| + \frac{C_0}{N} \sup_{0 \leq t \leq T} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right| + C_0 \varepsilon_N,$$

where \hat{x}_i is given by (32) and \bar{x}_i is given by (34). Here, $\varepsilon_N = \varepsilon_{1,N} + \varepsilon_{2,N} + \varepsilon_{3,N}$.

Proof. According to (14) and (32), it holds that

$$\left\{ \begin{aligned} d(\hat{x}_i^{(N)} - z_i) &= \left[A(\hat{x}_i^{(N)} - z_i) + \frac{B}{N} \sum_{j=1}^N g_{ij}^N \hat{u}_j - B \int_0^1 g(I_i^*, \beta) \mathbb{E} \bar{u}_\beta d\beta \right. \\ &\quad \left. + \frac{D}{N} \sum_{j=1}^N g_{ij}^N \hat{x}_j^{(N)} - D \int_0^1 g(I_i^*, \beta) z_\beta d\beta \right] dt + \frac{1}{N} \sum_{j=1}^N g_{ij}^N \sigma dw_j, \\ \hat{x}_i^{(N)}(0) - z_i(0) &= \frac{1}{N} \sum_{j=1}^N g_{ij}^N \xi_j - \int_0^1 g(I_i^*, \beta) m_\beta^x d\beta. \end{aligned} \right. \quad (37)$$

In the following estimates, we use C to denote a generic constant that does not depend on N and may vary from line to line. Recalling the control laws \hat{u}_i in (31), \bar{u}_i in (33) and \bar{u}_α in (10), at time t , we have

$$\begin{aligned} & \left| \frac{B}{N} \sum_{j=1}^N g_{ij}^N \hat{u}_j - B \int_0^1 g(I_i^*, \beta) \mathbb{E} \bar{u}_\beta d\beta \right| \\ &= \left| \frac{BR^{-1}B^\top}{N} \sum_{j=1}^N g_{ij}^N (\Pi \hat{x}_j + S_{I_j^*}) - BR^{-1}B^\top \int_0^1 g(I_i^*, \beta) (\Pi \mathbb{E} \bar{x}_\beta + S_\beta) d\beta \right| \\ &\leq C |\hat{x}_i^{(N)} - z_i| + C \left| \sum_{j=1}^N \left(\frac{g_{ij}^N}{N} - \int_{\beta \in I_j} g(I_i^*, \beta) d\beta \right) S_{I_j^*} \right| + C \left| \int_0^1 g(I_i^*, \beta) (S_\beta^N - S_\beta) d\beta \right| \\ &\leq C |\hat{x}_i^{(N)} - z_i| + C \varepsilon_N, \end{aligned}$$

where $S_\alpha^N = \sum_{j=1}^N \mathbb{1}_{I_j}(\alpha) S_{I_j^*}$. Here, the last inequality is due to the uniform boundedness of $S_{I_j^*}$ and $\sup_{0 \leq t \leq T} |S_\alpha^N - S_\alpha| \leq \sup_{0 \leq t \leq T} C |z_\alpha^N - z_\alpha|$ resulting from (14). Moreover, we also have

$$\begin{aligned} & \left| \frac{1}{N} \sum_{j=1}^N g_{ij}^N \xi_j - \int_0^1 g(I_i^*, \beta) m_\beta^x d\beta \right| \leq \frac{C}{N} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right| + C \varepsilon_N, \\ & \left| \frac{D}{N} \sum_{j=1}^N g_{ij}^N \hat{x}_j^{(N)} - D \int_0^1 g(I_i^*, \beta) z_\beta d\beta \right| \leq \frac{C}{N} \sum_{j=1}^N |\hat{x}_j^{(N)} - z_j| + C \varepsilon_N. \end{aligned}$$

Hence, by (37), for all $s \in [0, T]$, we obtain

$$\begin{aligned} |\hat{x}_i^{(N)}(s) - z_i(s)| &\leq C \int_0^s \left(|\hat{x}_i^{(N)} - z_i| + \frac{1}{N} \sum_{j=1}^N |\hat{x}_j^{(N)} - z_j| \right) dt + C \varepsilon_N \\ &\quad + \frac{C}{N} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right| + \frac{C}{N} \left| \int_0^s \sum_{j=1}^N \sigma dw_j \right|. \end{aligned} \quad (38)$$

Adding up the above N equations and using Gronwall's inequality, we derive

$$\frac{1}{N} \sum_{i=1}^N |\hat{x}_i^{(N)}(s) - z_i(s)| \leq \frac{C}{N} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right| + \frac{C}{N} \left| \int_0^s \sum_{j=1}^N \sigma dw_j \right| + C\varepsilon_N. \quad (39)$$

Combining (38) with (39), we get (35). Similarly, (36) can be proved. \square

For the risk-sensitive cost functionals (3) and (5), we need some exponentiated estimates of the difference between the processes in (32) and (34). For all the remaining part of this section, we take a sufficiently large \hat{N} and consider $N \geq \hat{N}$.

Lemma 5.3. Under (H1)–(H7), for any given deterministic positive function $\Theta = f(N)$, $N \in \mathbb{N}$, satisfying

$$f(N) = o\left(1/\left(\frac{1}{N} + \varepsilon_N^2\right)\right), \quad (40)$$

where ε_N is given by Lemma 5.2, we have

$$\sup_{1 \leq i \leq N} \mathbb{E}\left[e^{\Theta \int_0^T |\hat{x}_i - \bar{x}_i|^2 dt}\right] = 1 + o(1), \quad \sup_{1 \leq i \leq N} \mathbb{E}\left[e^{\Theta \int_0^T |\hat{x}_i^{(N)} - z_i|^2 dt}\right] = 1 + o(1). \quad (41)$$

Proof. By virtue of Lemma 5.2, for all $t \in [0, T]$, it holds that

$$|\hat{x}_i(t) - \bar{x}_i(t)|^2 \leq \frac{3C_0^2}{N^2} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right|^2 + \frac{3C_0^2}{N^2} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right| \right)^2 + 3C_0^2 \varepsilon_N^2.$$

In the following estimates, we use C to denote a generic constant that does not depend on N and may vary from line to line. Thus, it follows from Hölder's inequality that

$$\begin{aligned} & \mathbb{E} \left[\exp \left\{ \Theta T \left(\sup_{0 \leq t \leq T} |\hat{x}_i(t) - \bar{x}_i(t)|^2 \right) \right\} \right] \\ & \leq \left(\mathbb{E} \left[\exp \left\{ \frac{C\Theta}{N^2} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right|^2 \right\} \right] \right)^{\frac{1}{3}} \left(e^{C\Theta \varepsilon_N^2} \right)^{\frac{1}{3}} \\ & \quad \times \left(\mathbb{E} \left[\exp \left\{ \frac{C\Theta}{N^2} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right|^2 \right) \right\} \right] \right)^{\frac{1}{3}}. \end{aligned} \quad (42)$$

For the first term of the right-hand side of (42), using the independence of ξ_i as stated in (H1), we get

$$\mathbb{E} \left[\frac{C\Theta}{N^2} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right|^2 \right] = \frac{C\Theta}{N^2} \sum_{j=1}^N \mathbb{E} |\xi_j - \mathbb{E}\xi_j|^2 \leq C\Theta \left(\frac{1}{N} + \varepsilon_N^2 \right) \xrightarrow{N \rightarrow \infty} 0,$$

which implies

$$\lim_{N \rightarrow \infty} \frac{C\Theta}{N^2} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right|^2 = 0 \quad \text{in probability.} \quad (43)$$

Notice that $\xi_j, 1 \leq j \leq N$, are independent random variables taking values in a compact set S_0^x . Letting $Y_N = \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j)$, then by the integration by parts formula and Hoeffding's inequality, for all $\bar{K} > 1$, we have

$$\begin{aligned} & \mathbb{E} \left[e^{\frac{C\Theta}{N^2} |Y_N|^2} \mathbb{1}_{\left\{ e^{\frac{C\Theta}{N^2} |Y_N|^2} \geq \bar{K} \right\}} \right] \\ & \leq \int_{\bar{K}}^{\infty} \mathbb{P}(e^{\frac{C\Theta}{N^2} |Y_N|^2} > y) dy = \int_{\bar{K}}^{\infty} \mathbb{P}\left(|Y_N| > \sqrt{\frac{N^2 \ln y}{C\Theta}}\right) dy \\ & \leq \int_{\bar{K}}^{\infty} 2 \exp\left(-\frac{N^2 \ln y}{CN\Theta}\right) dy = \int_{\bar{K}}^{\infty} 2y^{-\frac{N}{C\Theta}} dy = \frac{2\bar{K}^{1-\frac{N}{C\Theta}}}{\frac{N}{C\Theta} - 1}, \end{aligned} \quad (44)$$

which tends to 0 as $\bar{K} \rightarrow \infty$, uniformly with respect to N , implying that the sequence $\{e^{\frac{C\Theta}{N^2}|Y_N|^2}\}_{N \geq \hat{N}}$ is uniformly integrable. Combining (44) with (43), it follows from Vitali convergence theorem (see [15, pp. 62-63]) that

$$\lim_{N \rightarrow \infty} \mathbb{E} \left[\exp \left\{ \frac{C\Theta}{N^2} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right|^2 \right\} \right] = 1. \quad (45)$$

For the second term of the right-hand side of (42), we have

$$\lim_{N \rightarrow \infty} e^{C\Theta \varepsilon_N^2} \leq \lim_{N \rightarrow \infty} e^{C\Theta(\frac{1}{N} + \varepsilon_N^2)} = 1. \quad (46)$$

For the last term of the right-hand side of (42), Since w_i , $1 \leq i \leq N$, are independent standard Brownian motions, $\int_0^t \sum_{j=1}^N \sigma dw_j$ has its distribution equal to that of $\sqrt{N} \int_0^t \sigma dw$, where w is a standard Brownian motion. Then, recalling condition (40), it follows from the monotonicity of e^x and Doob's maximal inequality for submartingales that, for sufficiently large $N \geq \hat{N}$,

$$\begin{aligned} \mathbb{E} \left[\exp \left\{ \frac{2C\Theta}{N^2} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right| \right)^2 \right\} \right] &= \mathbb{E} \left[\exp \left\{ \frac{2C\Theta}{N} \sup_{0 \leq t \leq T} \left| \int_0^t \sigma dw \right|^2 \right\} \right] \\ &\leq 4\mathbb{E} \left[\exp \left\{ \frac{2C\Theta}{N} \left| \int_0^T \sigma dw \right|^2 \right\} \right] \\ &= 4 \left[\det \left(I_n - \frac{4C\Theta}{N} \int_0^T \sigma \sigma^\top dt \right) \right]^{-\frac{1}{2}} \\ &= 4 + o(1), \end{aligned}$$

which implies that the sequence $\{e^{\frac{C\Theta}{N^2}(\sup_{0 \leq t \leq T} |\int_0^t \sum_{j=1}^N \sigma dw_j|)^2}\}_{N \geq \hat{N}}$ is uniformly integrable. Moreover, by Burkholder-Davis-Gundy inequality, we have

$$\mathbb{E} \left[\frac{C\Theta}{N^2} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right| \right)^2 \right] \leq \frac{C\Theta}{N} \int_0^T |\sigma|^2 dt \xrightarrow{N \rightarrow \infty} 0,$$

which implies that

$$\lim_{N \rightarrow \infty} \frac{C\Theta}{N^2} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right| \right)^2 = 0 \quad \text{in probability.}$$

Then, using Vitali convergence theorem again, we have

$$\lim_{N \rightarrow \infty} \mathbb{E} \left[\exp \left\{ \frac{C\Theta}{N^2} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right| \right)^2 \right\} \right] = 1. \quad (47)$$

Combining (42) with (45)–(47), it holds that

$$\mathbb{E} \left[e^{\Theta \int_0^T |\hat{x}_i - \bar{x}_i|^2 dt} \right] \leq \mathbb{E} \left[\exp \left\{ \Theta T \left(\sup_{0 \leq t \leq T} |\hat{x}_i - \bar{x}_i|^2 \right) \right\} \right] = 1 + o(1),$$

uniformly with respect to i . Then we obtain the first equation in (41). The proof of the second one is similar and omitted here. \square

For the processes (x, y, u) on $[0, T]$, define

$$\Lambda_T(x, y, u) = \int_0^T (\|x(t) - \Gamma y(t)\|_Q^2 + \|u(t)\|_R^2) dt + \|x(T) - \Gamma_f y(T)\|_{Q_f}^2, \quad (48)$$

and we further obtain the following estimate.

Lemma 5.4. Under (H1)–(H7), for all deterministic positive function $\Theta = f(N)$, $N \in \mathbb{N}$, satisfying

$$f(N) = o\left(1/\left(\sqrt{\frac{1}{N} + \varepsilon_N^2}\right)\right), \quad (49)$$

we have

$$\sup_{1 \leq i \leq N} \mathbb{E} \left[\exp \left\{ \Theta \left| \Lambda_T(\hat{x}_i, \hat{x}_i^{(N)}, \hat{u}_i) - \Lambda_T(\bar{x}_i, z_i, \bar{u}_i) \right| \right\} \right] = 1 + o(1). \quad (50)$$

Proof. Recalling (48), we have

$$\begin{aligned} & \left| \Lambda_T(\hat{x}_i, \hat{x}_i^{(N)}, \hat{u}_i) - \Lambda_T(\bar{x}_i, z_i, \bar{u}_i) \right| \\ & \leq \int_0^T \left\{ \left| \|\hat{x}_i - \Gamma \hat{x}_i^{(N)}\|_Q^2 - \|\bar{x}_i - \Gamma z_i\|_Q^2 \right| + \left| \|\hat{u}_i\|_R^2 - \|\bar{u}_i\|_R^2 \right| \right\} dt \\ & \quad + \left| \|\hat{x}_i(T) - \Gamma_f \hat{x}_i^{(N)}(T)\|_{Q_f}^2 - \|\bar{x}_i(T) - \Gamma_f z_i(T)\|_{Q_f}^2 \right|. \end{aligned}$$

Using Hölder's inequality, we get

$$\begin{aligned} & \mathbb{E} \left[\exp \left\{ \Theta \left| \Lambda_T(\hat{x}_i, \hat{x}_i^{(N)}, \hat{u}_i) - \Lambda_T(\bar{x}_i, z_i, \bar{u}_i) \right| \right\} \right] \\ & \leq \left(\mathbb{E} \left[e^{3\Theta \int_0^T \left| \|\hat{x}_i - \Gamma \hat{x}_i^{(N)}\|_Q^2 - \|\bar{x}_i - \Gamma z_i\|_Q^2 \right| dt} \right] \right)^{\frac{1}{3}} \left(\mathbb{E} \left[e^{3\Theta \int_0^T \left| \|\hat{u}_i\|_R^2 - \|\bar{u}_i\|_R^2 \right| dt} \right] \right)^{\frac{1}{3}} \\ & \quad \times \left(\mathbb{E} \left[e^{3\Theta \left| \|\hat{x}_i(T) - \Gamma_f \hat{x}_i^{(N)}(T)\|_{Q_f}^2 - \|\bar{x}_i(T) - \Gamma_f z_i(T)\|_{Q_f}^2 \right|} \right] \right)^{\frac{1}{3}}. \end{aligned} \quad (51)$$

Noticing $\langle Qx, x \rangle - \langle Qy, y \rangle = \langle Q(x - y), x - y \rangle + 2\langle Q(x - y), y \rangle$ and using Hölder's inequality, we have

$$\begin{aligned} & \mathbb{E} \left[e^{3\Theta \int_0^T \left| \|\hat{x}_i - \Gamma \hat{x}_i^{(N)}\|_Q^2 - \|\bar{x}_i - \Gamma z_i\|_Q^2 \right| dt} \right] \\ & \leq \mathbb{E} \left[e^{6\Theta \int_0^T \left\{ \|\hat{x}_i - \bar{x}_i\|_Q^2 + \|\hat{x}_i^{(N)} - z_i\|_{\Gamma^\top Q \Gamma}^2 + \langle Q[\hat{x}_i - \bar{x}_i - \Gamma(\hat{x}_i^{(N)} - z_i)], \bar{x}_i - \Gamma z_i \rangle \right\} dt} \right] \\ & \leq \left\{ \mathbb{E} \left[e^{C\Theta \int_0^T \|\hat{x}_i - \bar{x}_i\|^2 dt} \right] \right\}^{\frac{1}{4}} \left\{ \mathbb{E} \left[e^{C\Theta \int_0^T \left| \langle \hat{x}_i - \bar{x}_i, \bar{x}_i - \Gamma z_i \rangle \right| dt} \right] \right\}^{\frac{1}{4}} \\ & \quad \times \left\{ \mathbb{E} \left[e^{C\Theta \int_0^T \|\hat{x}_i^{(N)} - z_i\|^2 dt} \right] \right\}^{\frac{1}{4}} \left\{ \mathbb{E} \left[e^{C\Theta \int_0^T \left| \langle \hat{x}_i^{(N)} - z_i, \bar{x}_i - \Gamma z_i \rangle \right| dt} \right] \right\}^{\frac{1}{4}}, \end{aligned} \quad (52)$$

where $C > 0$ is a generic constant that does not depend on N and may vary from line to line. By virtue of Lemma 5.3, we know that

$$\left\{ \mathbb{E} \left[e^{C\Theta \int_0^T \|\hat{x}_i - \bar{x}_i\|^2 dt} \right] \right\}^{\frac{1}{4}} \left\{ \mathbb{E} \left[e^{C\Theta \int_0^T \|\hat{x}_i^{(N)} - z_i\|^2 dt} \right] \right\}^{\frac{1}{4}} = 1 + o(1). \quad (53)$$

Next, we analyze the other two factors in (52). Recalling the fundamental solution matrices $\Psi_z(t, s)$, we get

$$\bar{x}_i(t) = \Psi_z(t, 0)\xi_i - \int_0^t \Psi_z(t, s)(BR^{-1}B^\top S_{I_i^*} - Dz_i)ds + \int_0^t \Psi_z(t, s)\sigma dw_i.$$

By the GMFG equation system (14) and Theorem 4.1, we know that $(z_i, S_{I_i^*}) \in C([0, T]; \mathbb{R}^n) \times C([0, T]; \mathbb{R}^n)$, which implies that $(z_i, S_{I_i^*})$ is uniformly bounded with respect to all i . Since the initial state ξ_i takes value in a compact set S_0^x and $\Psi_z(t, s) = \Psi_z(t, 0)\Psi_z(0, s)$, we have

$$\sup_{0 \leq t \leq T} |\bar{x}_i - \Gamma z_i| \leq C + \sup_{0 \leq t \leq T} C \left| \int_0^t \Psi_z(0, s)\sigma dw_i \right|.$$

Using Hölder's inequality and a method similar to the proof of Lemma 5.3, taking $\tilde{\varepsilon} = \sqrt{\frac{1}{N} + \varepsilon_N^2}$, then we obtain

$$\mathbb{E} \left[\exp \left(C\Theta T \cdot \sup_{0 \leq t \leq T} |\langle \hat{x}_i - \bar{x}_i, \bar{x}_i - \Gamma z_i \rangle| \right) \right]$$

$$\begin{aligned}
&\leq \mathbb{E} \left[\exp \left(\frac{C_2 \Theta T}{2\tilde{\epsilon}} \cdot \sup_{0 \leq t \leq T} |\hat{x}_i - \bar{x}|^2 + \frac{\tilde{\epsilon} C \Theta T}{2} \cdot \sup_{0 \leq t \leq T} |\bar{x}_i - \Gamma z_i|^2 \right) \right] \\
&\leq \left\{ \mathbb{E} \left[\exp \left(\frac{C \Theta T}{\tilde{\epsilon}} \cdot \sup_{0 \leq t \leq T} |\hat{x}_i - \bar{x}_i|^2 \right) \right] \right\}^{\frac{1}{2}} \left\{ \mathbb{E} \left[\exp \left(\tilde{\epsilon} C \Theta T \cdot \sup_{0 \leq t \leq T} |\bar{x}_i - \Gamma z_i|^2 \right) \right] \right\}^{\frac{1}{2}} \\
&\leq [1 + o(1)] \left\{ \mathbb{E} \left[\exp \left\{ \tilde{\epsilon} C \Theta T \left[C^2 + \left(\sup_{0 \leq t \leq T} C \left| \int_0^t \Psi_z(0, s) \sigma dw_i \right| \right)^2 \right] \right\} \right] \right\}^{\frac{1}{2}} \\
&\leq [1 + o(1)] e^{\tilde{\epsilon} C \Theta T} \left\{ \mathbb{E} \left[\exp \left\{ \tilde{\epsilon} C \Theta T \left(\sup_{0 \leq t \leq T} \left| \int_0^t \Psi_z(0, s) \sigma dw_i \right| \right)^2 \right\} \right] \right\}^{\frac{1}{4}} \\
&= 1 + o(1),
\end{aligned}$$

where the second to last line is due to Vitali convergence theorem and

$$\begin{aligned}
&\lim_{N \rightarrow \infty} \tilde{\epsilon} C \Theta T \left(\sup_{0 \leq t \leq T} \left| \int_0^t \Psi_z(0, s) \sigma dw_i \right| \right)^2 = 0, \quad \text{in probability,} \\
&\sup_{N \geq \tilde{N}} \mathbb{E} \left[\exp \left\{ 2\tilde{\epsilon} C \Theta T \left(\sup_{0 \leq t \leq T} \left| \int_0^t \Psi_z(0, s) \sigma dw_i \right| \right)^2 \right\} \right] \leq 4 + o(1).
\end{aligned}$$

Similarly, we also have

$$\mathbb{E} \left[\exp \left(C \Theta T \cdot \sup_{0 \leq t \leq T} |\langle \hat{x}_i^{(N)} - z_i, \bar{x}_i - \Gamma z_i \rangle| \right) \right] = 1 + o(1).$$

Making use of the above estimates, we further obtain

$$\begin{aligned}
&\left\{ \mathbb{E} \left[e^{C \Theta \int_0^T |\langle \hat{x}_i - \bar{x}_i, \bar{x}_i - \Gamma z_i \rangle| dt} \right] \right\}^{\frac{1}{4}} \left\{ \mathbb{E} \left[e^{C \Theta \int_0^T |\langle \hat{x}_i^{(N)} - z_i, \bar{x}_i - \Gamma z_i \rangle| dt} \right] \right\}^{\frac{1}{4}} \\
&\leq \left\{ \mathbb{E} \left[\exp \left\{ C \Theta T \left(\sup_{0 \leq t \leq T} |\langle \hat{x}_i - \bar{x}_i, \bar{x}_i - \Gamma z_i \rangle| \right) \right\} \right] \right\}^{\frac{1}{4}} \\
&\quad \times \left\{ \mathbb{E} \left[\exp \left\{ C \Theta T \left(\sup_{0 \leq t \leq T} |\langle \hat{x}_i^{(N)} - z_i, \bar{x}_i - \Gamma z_i \rangle| \right) \right\} \right] \right\}^{\frac{1}{4}} \\
&= 1 + o(1).
\end{aligned} \tag{54}$$

Combining (53) with (54), we have

$$\mathbb{E} \left[e^{3\Theta \int_0^T \|\hat{x}_i - \Gamma \hat{x}_i^{(N)}\|_Q^2 - \|\bar{x}_i - \Gamma z_i\|_Q^2 dt} \right] = 1 + o(1). \tag{55}$$

Applying a similar method as used in the proof of (55), we also obtain

$$\begin{aligned}
&\mathbb{E} \left[e^{3\Theta \int_0^T \|\hat{u}_i\|_R^2 - \|\bar{u}_i\|_R^2 dt} \right] = 1 + o(1), \\
&\mathbb{E} \left[e^{3\Theta \left[\|\hat{x}_i(T) - \Gamma_f \hat{x}_i^{(N)}(T)\|_{Q_f}^2 - \|\bar{x}_i(T) - \Gamma_f z_i(T)\|_{Q_f}^2 \right]} \right] = 1 + o(1).
\end{aligned} \tag{56}$$

Combining (51) with (55)–(56), we obtain (50). \square

Suppose that agent \mathcal{A}_i chooses an alternative feedback control law $u_i \in \mathcal{U}_c^i$, while the other agents \mathcal{A}_j , $j \neq i$, still adopt the strategies \hat{u}_j in (31). The resulting closed-loop state processes are given by

$$\begin{cases} dx_i = (Ax_i + Bu_i + Dx_i^{(N)})dt + \sigma dw_i, & x_i(0) = \xi_i, \\ dx_j = [Ax_j - BR^{-1}B^\top(\Pi x_j + S I_j^*) + Dx_j^{(N)}]dt + \sigma dw_j, & x_j(0) = \xi_j, \quad j \neq i, \end{cases} \tag{57}$$

where $x_i^{(N)} = \frac{1}{N} \sum_{l=1}^N g_{il}^N x_l$. The cost functional of agent \mathcal{A}_i is

$$\mathcal{J}_i(u_i, \hat{u}_{-i}) = \mathbb{E} \left[\exp \left(\gamma \left\{ \int_0^T [\|x_i - \Gamma x_i^{(N)}\|_Q^2 + \|u_i\|_R^2] dt + \|x_i(T) - \Gamma_f x_i^{(N)}(T)\|_{Q_f}^2 \right\} \right) \right].$$

Let $u_i^{cl}(t, \omega)$, as a stochastic process adapted to \mathcal{F}_t , be the closed-loop control inputs generated by the above control law u_i . Based on (4), we introduce the auxiliary mean-field limit dynamics for each agent:

$$\begin{cases} dx_i^\infty = (Ax_i^\infty + Bu_i^{cl} + Dz_i)dt + \sigma dw_i, & x_i^\infty(0) = \xi_i, \\ dx_j^\infty = [Ax_j^\infty - BR^{-1}B^\top(\Pi x_j^\infty + S_{I_j^*}) + Dz_j]dt + \sigma dw_j, & x_j^\infty(0) = \xi_j, \quad j \neq i, \end{cases} \quad (58)$$

where $z_i = z_{I_i^*}$ is specified by (14) and u_i^{cl} is the closed-loop control inputs generated by the control law u_i in (57). The use of the form u_i^{cl} is for convenience of further performance estimate (see the subsequent proof of Theorem 5.6). Then, by the method of proving Lemma 5.2, we obtain the following estimates.

Lemma 5.5. Under (H1)–(H7), for (57) and (58), there exists a constant \bar{C} such that

$$\sup_{1 \leq i \leq N} \sup_{0 \leq t \leq T} |x_i^{(N)} - z_i| \leq \bar{C} \left(\frac{1}{N} + \varepsilon_N \right) + \frac{\bar{C}}{N} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right| \quad (59)$$

$$+ \frac{\bar{C}}{N} \int_0^T |u_i^{cl}| dt + \frac{\bar{C}}{N} \sup_{0 \leq t \leq T} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right|,$$

$$\sup_{1 \leq i \leq N} \sup_{0 \leq t \leq T} |x_i - x_i^\infty| \leq \bar{C} \left(\frac{1}{N} + \varepsilon_N \right) + \frac{\bar{C}}{N} \left| \sum_{j=1}^N (\xi_j - \mathbb{E}\xi_j) \right| \quad (60)$$

$$+ \frac{\bar{C}}{N} \int_0^T |u_i^{cl}| dt + \frac{\bar{C}}{N} \sup_{0 \leq t \leq T} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right|.$$

Proof. By (14) and (57), it holds that

$$\begin{cases} dx_i^{(N)} - z_i = \left[A(x_i^{(N)} - z_i) + \frac{B}{N} g_{ii}^N u_i^{cl} - \frac{BR^{-1}B^\top}{N} \sum_{j \neq i} g_{ij}^N (\Pi x_j + S_{I_j^*}) + \frac{D}{N} \sum_{j=1}^N g_{ij}^N x_j^{(N)} \right. \\ \quad \left. - B \int_0^1 g(I_i^*, \beta) \mathbb{E} \bar{u}_\beta d\beta - D \int_0^1 g(I_i^*, \beta) z_\beta d\beta \right] dt + \frac{1}{N} \sum_{j=1}^N g_{ij}^N \sigma dw_j, \\ x_i^{(N)}(0) - z_i(0) = \frac{1}{N} \sum_{j=1}^N g_{ij}^N \xi_j - \int_0^1 g(I_i^*, \beta) m_\beta^x d\beta. \end{cases} \quad (61)$$

In the following estimates, we again use C to denote a generic constant that does not depend on N and may vary from line to line. Recalling the control laws \bar{u}_i in (33) and \bar{u}_α in (10), we have

$$\begin{aligned} & \int_0^s \left| \frac{B}{N} g_{ii}^N u_i^{cl} - \frac{BR^{-1}B^\top}{N} \sum_{j \neq i} g_{ij}^N (\Pi x_j + S_{I_j^*}) - B \int_0^1 g(I_i^*, \beta) \mathbb{E} \bar{u}_\beta d\beta \right| dt \\ & \leq \int_0^s \left| \frac{B}{N} g_{ii}^N u_i^{cl} + \frac{BR^{-1}B^\top}{N} g_{ii}^N (\Pi x_i + S_{I_i^*}) \right| dt \\ & \quad + \int_0^s \left| \frac{BR^{-1}B^\top}{N} \sum_{j=1}^N g_{ij}^N (\Pi x_j + S_{I_j^*}) - BR^{-1}B^\top \int_0^1 g(I_i^*, \beta) (\Pi \bar{x}_\beta + S_\beta) d\beta \right| dt \\ & \leq \int_0^s \left| \frac{B}{N} g_{ii}^N u_i^{cl} + \frac{BR^{-1}B^\top}{N} g_{ii}^N (\Pi x_i + S_{I_i^*}) \right| dt + C \int_0^s |x_i^{(N)} - z_i| dt + C \varepsilon_N^2 \\ & \leq \frac{C}{N} \int_0^s |u_i^{cl}| dt + C \int_0^s |x_i^{(N)} - z_i| dt + \frac{C}{N} \sup_{0 \leq t \leq s} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right| + C \left(\frac{1}{N} + \varepsilon_N \right), \end{aligned}$$

where the last inequality is due to $|x_i(s)| \leq C + C \int_0^s |u_i^{cl}| dt + C \left| \int_0^s \sum_{j=1}^N \sigma dw_j \right|$. Moreover, we also have

$$\begin{aligned} \left| \frac{1}{N} \sum_{j=1}^N g_{ij}^N \xi_j - \int_0^1 g(I_i^*, \beta) m_\beta^x d\beta \right| &\leq \frac{C}{N} \left| \sum_{j=1}^N (\xi_j - \mathbb{E} \xi_j) \right| + C \varepsilon_N, \\ \left| \frac{D}{N} \sum_{j=1}^N g_{ij}^N x_j^{(N)} - D \int_0^1 g(I_i^*, \beta) z_\beta d\beta \right| &\leq \frac{C}{N} \sum_{j=1}^N |x_j^{(N)} - z_j| + C \varepsilon_N. \end{aligned}$$

Thus, by (61), for all $s \in [0, T]$, we have

$$\begin{aligned} |x_i^{(N)}(s) - z_i(s)| &\leq C \int_0^s \left(|x_i^{(N)} - z_i| + \frac{1}{N} \sum_{j=1}^N |x_j^{(N)} - z_j| \right) dt + C \left(\frac{1}{N} + \varepsilon_N \right) \\ &\quad + \frac{C}{N} \left| \sum_{j=1}^N (\xi_j - \mathbb{E} \xi_j) \right| + \frac{C}{N} \int_0^s |u_i^{cl}| dt + \frac{C}{N} \sup_{0 \leq t \leq s} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right|. \end{aligned} \quad (62)$$

Adding up the above N equations and using Gronwall's inequality, we have

$$\begin{aligned} \frac{1}{N} \sum_{j=1}^N |x_j^{(N)}(s) - z_j(s)| &\leq C \left(\frac{1}{N} + \varepsilon_N \right) + \frac{C}{N} \left| \sum_{j=1}^N (\xi_j - \mathbb{E} \xi_j) \right| + \frac{C}{N} \int_0^t |u_i^{cl}| dt + \frac{C}{N} \sup_{0 \leq t \leq s} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right|. \end{aligned} \quad (63)$$

Combining (62) with (63), we get (59). Similarly, (60) can be proved. \square

Based on the above analysis, we now present the key result on performance analysis.

Theorem 5.6. Suppose (H1)–(H7) hold. The set of strategies $\hat{u} = (\hat{u}_1, \dots, \hat{u}_N)$ given by (31) is an ε -Nash equilibrium, where $\varepsilon = o(1)$ as $N \rightarrow \infty$.

According to Definition 5.1 of the ε -Nash equilibrium, it suffices to show

$$\mathcal{J}_i(\hat{u}_i, \hat{u}_{-i}) \leq \inf_{u_i \in \mathcal{U}_i^c} \mathcal{J}_i(u_i, \hat{u}_{-i}) + \varepsilon.$$

Since the classical method of proving the ε -Nash equilibrium via L^2 error estimates alone is inadequate, a new method is applied subsequently. We will identify sufficiently close upper and lower bounds of $\inf_{u_i \in \mathcal{U}_i^c} \mathcal{J}_i(u_i, \hat{u}_{-i})$. Subsequently, we can show that the set of decentralized strategies \hat{u} is an ε -Nash equilibrium.

Proof of Theorem 5.6. Step 1. Provided that we can show $J_i(\hat{u}_i, \hat{u}_{-i}) \leq \tilde{C}$ for some fixed constant \tilde{C} independent of $N \geq \hat{N}$, then it is sufficient to consider all alternative control laws $u_i \in \mathcal{U}_i^c$ that satisfy

$$\mathcal{J}_i(u_i, \hat{u}_{-i}) \leq \tilde{C}. \quad (64)$$

Recalling the definition of $\Lambda_T(x, y, u)$ in (48) and using Hölder's inequality, we obtain

$$\begin{aligned} \mathcal{J}_i(\hat{u}_i, \hat{u}_{-i}) &= \mathbb{E} \left[e^{\gamma \Lambda_T(\hat{x}_i, \hat{x}_i^{(N)}, \hat{u}_i)} \right] \\ &\leq \left(\mathbb{E} \left[e^{\gamma(1+\delta) \Lambda_T(\bar{x}_i, z_i, \bar{u}_i)} \right] \right)^{\frac{1}{1+\delta}} \left(\mathbb{E} \left[e^{\frac{\gamma(1+\delta)}{\delta} [\Lambda_T(\hat{x}_i, \hat{x}_i^{(N)}, \hat{u}_i) - \Lambda_T(\bar{x}_i, z_i, \bar{u}_i)]} \right] \right)^{\frac{\delta}{1+\delta}}, \end{aligned}$$

where (\bar{x}_i, \bar{u}_i) is specified by (33)–(34) and $z_i = z_{I_i^*}$ is specified by (14). Selecting $\delta_N = \frac{\gamma \sqrt{\frac{1}{N} + \varepsilon_N^2}}{1 - \gamma \sqrt{\frac{1}{N} + \varepsilon_N^2}}$, we see that $\delta_N \rightarrow 0$ as $N \rightarrow \infty$. We will use the fact that $(\mathbb{E}|X|^{1+\delta})^{\frac{1}{1+\delta}}$ approaches $\mathbb{E}|X|$ as $\delta \rightarrow 0$, if there exists a positive δ_0 such that $\mathbb{E}|X|^{1+\delta_0} < \infty$. By Theorem 3.1, we have

$$\mathbb{E} \left[\exp \left\{ \gamma \Lambda_T(\bar{x}_i, z_i, \bar{u}_i) \right\} \right] = \mathbb{E} \left[\exp \left\{ \gamma \left\{ \xi_i^\top \Pi(0) \xi_i + 2 \xi_i^\top S_{I_i^*}(0) + r_{I_i^*}(0) \right\} \right\} \right].$$

In conjunction with (H1) and the uniform boundedness of $(\Pi, S_{I_i^*}, r_{I_i^*})$, it follows that $\mathbb{E}[e^{\gamma(1+\delta_0)\Lambda_T(\bar{x}_i, z_i, \bar{u}_i)}] < \infty$ for some positive constant δ_0 . Thus

$$\lim_{N \rightarrow \infty} \left(\mathbb{E} \left[\exp \left\{ \gamma(1 + \delta_N) \Lambda_T(\bar{x}_i, z_i, \bar{u}_i) \right\} \right] \right)^{\frac{1}{1+\delta_N}} = \mathbb{E} \left[\exp \left(\gamma \Lambda_T(\bar{x}_i, z_i, \bar{u}_i) \right) \right]. \quad (65)$$

Now we take $\Theta = \frac{\gamma(1+\delta_N)}{\delta_N} = \frac{1}{\sqrt{\frac{1}{N} + \varepsilon_N^2}}$, and we can verify that condition (49) holds. By Lemma 5.4, we get

$$\mathbb{E} \left[\exp \left\{ \frac{\gamma(1 + \delta_N)}{\delta_N} \left| \Lambda_T(\hat{x}_i, \hat{x}_i^{(N)}, \hat{u}_i) - \Lambda_T(\bar{x}_i, z_i, \bar{u}_i) \right| \right\} \right] = 1 + o(1). \quad (66)$$

Then combining (65) with (66), it holds that

$$\mathcal{J}_i(\hat{u}_i, \hat{u}_{-i}) \leq J_i(\bar{u}_i)[1 + o(1)], \quad (67)$$

where \bar{u}_i is given by (33), and $o(1) \rightarrow 0$ as $N \rightarrow \infty$. Hence, there exists a fixed constant \tilde{C} such that $\sup_{1 \leq i \leq N} \mathcal{J}_i(\hat{u}_i, \hat{u}_{-i}) \leq \tilde{C}$ indeed holds for all $N \geq \hat{N}$. In view of (H3) and (64), we have

$$\mathbb{E}[e^{\gamma \int_0^T \|u_i\|_R^2 dt}] \leq \mathcal{J}_i(u_i, \hat{u}_{-i}) \leq \tilde{C}, \quad (68)$$

which implies that, for all sufficiently large N , there exists a constant \tilde{C}_1 (depending on \tilde{C} and the uniform upper bound of R),

$$\mathbb{E} \int_0^T |u_i|^2 dt \leq \frac{\tilde{C}_1}{\gamma}. \quad (69)$$

Step 2. For $u_i \in \mathcal{U}_i^i$ satisfying $\mathcal{J}_i(u_i, \hat{u}_{-i}) \leq \tilde{C}$, we aim to find a lower bound of $\mathcal{J}_i(u_i^{cl}, \hat{u}_{-i})$, where u_i^{cl} appears in (58) and now additionally satisfies (68)–(69).

Let X and Y be two random variables. For any given positive constant c , using Hölder's inequality, we have $\mathbb{E}[e^{cY}] \leq (\mathbb{E}[e^{\frac{c(1+\delta')}{\delta'}(Y-X)}])^{\frac{\delta'}{1+\delta'}} (\mathbb{E}[e^{c(1+\delta')X}])^{\frac{1}{1+\delta'}}$ for every $\delta' > 0$. Taking $c = \frac{1}{1+\delta'}$, it holds that $\mathbb{E}[e^X] \geq (\mathbb{E}[e^{\frac{1}{1+\delta'}Y}])^{1+\delta'} (\mathbb{E}[e^{\frac{1}{\delta'}(Y-X)}])^{-\delta'}$. Based on this inequality, we have

$$\mathbb{E}[e^{\gamma \Lambda_T(x_i, x_i^{(N)}, u_i^{cl})}] \geq \frac{(\mathbb{E}[e^{\frac{\gamma}{1+\delta'} \Lambda_T(x_i^\infty, z_i, u_i^{cl})}])^{1+\delta'}}{(\mathbb{E}[e^{\frac{\gamma}{\delta'} [\Lambda_T(x_i^\infty, z_i, u_i^{cl}) - \Lambda_T(x_i, x_i^{(N)}, u_i^{cl})]}])^{\delta'}}, \quad (70)$$

where the control u_i^{cl} is the same as in (58). Note that the alternative control law u_i satisfies (68)–(69). Then in view of Lemma 5.5, the estimate in Lemma 5.4 after replacing \hat{u}_i by u_i^{cl} remains valid (see (77)), which guarantees that the numerator and the denominator in (70) are not simultaneously 0 or ∞ .

To estimate the numerator in (70), we introduce a new auxiliary control problem (ACP), which considers the further optimization of u_i^{cl} appearing in the numerator of (70).

Problem (ACP). Find $\bar{u}_i^{\delta'} \in \mathcal{U}_i^i$ to minimize

$$J_i^{\delta'}(u_i) = \mathbb{E} \left[\exp \left(\frac{\gamma}{1+\delta'} \left\{ \int_0^T [\|x_i - \Gamma z_i\|_Q^2 + \|u_i\|_R^2] dt + \|x_i(T) - \Gamma_f z_i(T)\|_{Q_f}^2 \right\} \right) \right],$$

for $\delta' > 0$, subject to

$$dx_i = (Ax_i + Bu_i + Dz_i)dt + \sigma dw_i, \quad x_i(0) = \xi_i,$$

where $z_i = z_{I_i^*}$ is specified by (14).

By the method of Section 3, we obtain the optimal control law

$$\bar{u}_i^{\delta'} = -R^{-1}B^\top \Pi^{\delta'} \bar{x}_i^{\delta'} - R^{-1}B^\top S_i^{\delta'},$$

where $\bar{x}_i^{\delta'}$ satisfies

$$d\bar{x}_i^{\delta'} = [(A - BR^{-1}B^\top \Pi^{\delta'})\bar{x}_i^{\delta'} - BR^{-1}B^\top S_i^{\delta'} + Dz_i]dt + \sigma dw_i, \quad \bar{x}_i^{\delta'}(0) = \xi_i.$$

Here, $\Pi^{\delta'}$ is given by the following Riccati equation

$$\dot{\Pi}^{\delta'} + \Pi^{\delta'} A + A^\top \Pi^{\delta'} - \Pi^{\delta'} \left(BR^{-1}B^\top - \frac{2\gamma}{1+\delta'} \sigma \sigma^\top \right) \Pi^{\delta'} + Q = 0, \quad \Pi^{\delta'}(T) = Q_f, \quad (71)$$

and $S_i^{\delta'}$ satisfies the ODE

$$\dot{S}_i^{\delta'} + \left(A^\top - \Pi^{\delta'} BR^{-1}B^\top + \frac{2\gamma}{1+\delta'} \Pi^{\delta'} \sigma \sigma^\top \right) S_i^{\delta'} - (Q\Gamma - \Pi^{\delta'} D)z_i = 0, \quad (72)$$

with $S_i^{\delta'}(T) = -Q_f \Gamma_f z_i(T)$. By (H4), we have $BR^{-1}B^\top - \frac{2\gamma}{1+\delta'} \sigma \sigma^\top \geq 0$ for each $t \in [0, T]$. Then Riccati equation (71) admits a unique solution $\Pi^{\delta'} \in C([0, T]; \mathbb{S}^n)$ under (H2)–(H4). Furthermore, ODE (72) also admits a unique solution $S_i^{\delta'} \in C([0, T]; \mathbb{R}^n)$.

The optimal cost is given by

$$J_i^{\delta'}(\bar{u}_i^{\delta'}) = \mathbb{E} \left[\exp \left(\frac{\gamma}{1+\delta'} \{ \xi_i^\top \Pi^{\delta'}(0) \xi_i + 2 \xi_i^\top S_i^{\delta'}(0) + r_i^{\delta'}(0) \} \right) \right], \quad (73)$$

where $r_i^{\delta'}$ satisfies

$$\dot{r}_i^{\delta'} - (S_i^{\delta'})^\top \left(BR^{-1}B^\top - \frac{2\gamma}{1+\delta'} \sigma \sigma^\top \right) S_i^{\delta'} + 2z_i^\top D^\top S_i^{\delta'} + z_i^\top \Gamma^\top Q \Gamma z_i + \text{Tr}(\sigma \sigma^\top \Pi^{\delta'}) = 0, \quad (74)$$

with $r_i^{\delta'}(T) = z_i^\top(T) \Gamma_f^\top Q_f \Gamma_f z_i(T)$. Under (H2)–(H4), ODE (74) admits a unique solution $r_i^{\delta'} \in C([0, T]; \mathbb{R})$.

We proceed to estimate the denominator on the right-hand side of (70). For (57)–(58), by Lemma 5.5 and Hölder's inequality, it holds that

$$\begin{aligned} & \mathbb{E} \left[\exp \left\{ \Theta T \left(\sup_{0 \leq t \leq T} |x_i(t) - x_i^\infty(t)|^2 \right) \right\} \right] \\ & \leq \left(\mathbb{E} \left[\exp \left\{ \frac{16\Theta T \bar{C}^2}{N^2} \left| \sum_{j=1}^N (\xi_j - \mathbb{E} \xi_j) \right|^2 \right\} \right] \right)^{\frac{1}{4}} \left(\mathbb{E} \left[\exp \left\{ \frac{16\Theta T \bar{C}^2}{N^2} \int_0^T |u_i^{cl}|^2 dt \right\} \right] \right)^{\frac{1}{4}} \\ & \quad \times \left(\mathbb{E} \left[\exp \left\{ \frac{16\Theta T C_0^2}{N^2} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \sum_{j=1}^N \sigma dw_j \right| \right)^2 \right\} \right] \right)^{\frac{1}{4}} \left(e^{16\Theta T \bar{C}^2 (\frac{1}{N} + \varepsilon_N^2)} \right)^{\frac{1}{4}}. \end{aligned} \quad (75)$$

Noticing (68)–(69), for all deterministic positive function $\Theta = f(N)$, $N \in \mathbb{N}$, satisfying $f(N) = o(1/\sqrt{\frac{1}{N} + \varepsilon_N^2})$, we have

$$\sup_{N \geq \bar{N}} \mathbb{E} \left[\exp \left\{ \frac{32\Theta T \bar{C}^2}{N^2} \int_0^T |u_i^{cl}|^2 dt \right\} \right] \leq \tilde{C}, \quad \frac{16\Theta T \bar{C}^2}{N^2} \int_0^T |u_i^{cl}|^2 dt \xrightarrow[N \rightarrow \infty]{\text{in probability}} 0,$$

which by Vitali convergence theorem implies that

$$\mathbb{E} \left[\exp \left\{ \frac{16\Theta T \bar{C}^2}{N^2} \int_0^T |u_i^{cl}|^2 dt \right\} \right] = 1 + o(1).$$

By the method of proving Lemma 5.3, we use (75) to obtain

$$\mathbb{E} [e^{\Theta \int_0^T |x_i - x_i^\infty|^2 dt}] = 1 + o(1), \quad \text{and} \quad \mathbb{E} [e^{\Theta \int_0^T |x_i^{(N)} - z_i|^2 dt}] = 1 + o(1). \quad (76)$$

By use of (76), we follow the proof of Lemma 5.4 to establish

$$\mathbb{E}\left[\exp\left\{\Theta\left|\Lambda_T(x_i^\infty, z_i, u_i^{cl}) - \Lambda_T(x_i, x_i^{(N)}, u_i^{cl})\right|\right\}\right] = 1 + o(1). \quad (77)$$

Now, we select $\delta'_N = \frac{1}{\gamma} \sqrt{\frac{1}{N} + \varepsilon_N^2}$ and $\Theta = \frac{\gamma}{\delta'_N} = \frac{1}{\sqrt{\frac{1}{N} + \varepsilon_N^2}}$, and it follows from (77) that

$$\left(\mathbb{E}\left[\exp\left\{\frac{\gamma}{\delta'_N}\left|\Lambda_T(x_i^\infty, z_i, u_i^{cl}) - \Lambda_T(x_i, x_i^{(N)}, u_i^{cl})\right|\right\}\right]\right)^{\delta'_N} = 1 + o(1). \quad (78)$$

Combining (70), (73) with (78), we have

$$\mathcal{J}_i(u_i^{cl}, \hat{u}_{-i}) = \mathbb{E}[e^{\gamma\Lambda_T(x_i, x_i^{(N)}, u_i^{cl})}] \geq \frac{[J_i^{\delta'_N}(\bar{u}_i^{\delta'_N})]^{1+\delta'_N}}{1+o(1)} = [J_i^{\delta'_N}(\bar{u}_i^{\delta'_N})]^{1+\delta'_N} [1+o(1)]. \quad (79)$$

Step 3. We proceed to compare the perturbation of the cost functionals. By the continuous dependence of solutions on parameters in ODEs, we have

$$\lim_{N \rightarrow \infty} (\Pi^{\delta'_N}, S_i^{\delta'_N}, r_i^{\delta'_N}) = (\Pi, S_{I_i^*}, r_{I_i^*}),$$

uniformly on $[0, T]$, which implies (recalling (12), (73))

$$\lim_{N \rightarrow \infty} J_i^{\delta'_N}(\bar{u}_i^{\delta'_N}) = J_i(\bar{u}_i).$$

Noticing the upper bound (67) and lower bound (79) of $\mathcal{J}_i(u_i^{cl}, \hat{u}_{-i})$, for all sufficiently large N , we have

$$J_i(\bar{u}_i)[1+o(1)] \leq \inf_{u_i \in \mathcal{U}_c^i} \mathcal{J}_i(u_i, \hat{u}_{-i}) \leq \mathcal{J}_i(\hat{u}_i, \hat{u}_{-i}) \leq J_i(\bar{u}_i)[1+o(1)],$$

where $J_i(\bar{u}_i) \times o(1) \rightarrow 0$ as $N \rightarrow \infty$, uniformly with respect to i . Then we have

$$\lim_{N \rightarrow \infty} \sup_{1 \leq i \leq N} \left| \inf_{u_i \in \mathcal{U}_c^i} \mathcal{J}_i(u_i, \hat{u}_{-i}) - \mathcal{J}_i(\hat{u}_i, \hat{u}_{-i}) \right| = 0, \quad (80)$$

which completes the proof of the theorem. \square

Remark 5.7. i) The exponentiated cost structure here makes the performance estimate more difficult than in LQG MFGs with risk-neutral costs for which standard L^2 error estimates are adequate [10, 26].

ii) If the initial states are independent Gaussian random variables with uniformly bounded second moments, we can still establish the ε -Nash equilibrium theorem. In this case, the compactness condition in (H1) is not fulfilled, but we can directly estimate (45) using the method in [36, pp. 39-42] instead of Hoeffding's inequality.

6 Numerical example

In this section, we provide a numerical example. We first compute the function z_α , which is specified by (14) or equivalently by (18). Here we will use (18) to facilitate the numerical method developed below. Inspired by [23], we decouple the forward-backward equation (18) by the following asymmetric operator Riccati equation

$$\begin{cases} \dot{\mathbf{P}} + \mathbf{P}(\mathbb{A} + [\mathbf{D}\mathbf{G}]) + \mathbb{A}^\top \mathbf{P} + [2\gamma \Pi \sigma \sigma^\top \mathbb{I}] \mathbf{P} - \mathbf{P}[\mathbf{B}\mathbf{R}^{-1}\mathbf{B}^\top \mathbf{G}] \mathbf{P} - [(\mathbf{Q}\mathbf{I} - \Pi \mathbf{D})\mathbb{I}] = 0, \\ \mathbf{P}(T) = -[\mathbf{Q}_f \mathbf{I}_f \mathbb{I}]. \end{cases} \quad (81)$$

In analogue to [23, Proposition 8], we can show that the operator Riccati equation (81) admits a unique mild solution (see [23, Lemma 1]) if $C_\Xi < 1$, where C_Ξ is defined by (4.1).

Next, we will derive the spectral decomposition of this operator Riccati equation, which allows us to approximate the operator Riccati equation by use of some finite dimensional matrix valued Riccati equations. By [23, Proposition 6], we have the following spectral composition of Riccati equation (81):

$$\mathbf{P}(t) = [P^\perp(t)\mathbb{I}] + \sum_{\ell \in \mathcal{I}_\lambda} [(P^\ell(t) - P^\perp(t))\mathbf{f}_\ell \mathbf{f}_\ell^\top].$$

where $\{\mathbf{f}_\ell\}_{\ell \in \mathcal{I}_\lambda}$ denotes the orthonormal eigenfunctions of \mathbf{G} ; λ_ℓ denotes the eigenvalue of \mathbf{G} corresponding to \mathbf{f}_ℓ ; and \mathcal{I}_λ is the index multiset for all the nonzero eigenvalues of \mathbf{G} . For further details on the eigenvalues, see Example 4.7.

Moreover, P^\perp and P^ℓ are respectively given by

$$\begin{cases} \dot{P}^\perp + P^\perp(A - BR^{-1}B^\top \Pi) + (A - BR^{-1}B^\top \Pi)^\top P^\perp + 2\gamma \Pi \sigma \sigma^\top P^\perp - (Q\Gamma - \Pi D) = 0, \\ P^\perp(T) = -Q_f \Gamma_f, \end{cases} \quad (82)$$

and

$$\begin{cases} \dot{P}^\ell + P^\ell(A - BR^{-1}B^\top \Pi + \lambda_\ell D) + (A - BR^{-1}B^\top \Pi + 2\gamma \Pi \sigma \sigma^\top)P^\ell \\ \quad - \lambda_\ell P^\ell BR^{-1}B^\top P^\ell - (Q\Gamma - \Pi D) = 0, \\ P^\ell(T) = -Q_f \Gamma_f. \end{cases} \quad (83)$$

For illustration, we now consider a sinusoidal graphon as follows

$$\mathbf{G}(\alpha, \beta) = \cos^2\left(\frac{\pi}{2}(\alpha - \beta)\right), \quad \alpha, \beta \in [0, 1], \quad (84)$$

which is illustrated in Figure 1a. For this sinusoidal graphon \mathbf{G} , we can show that the normalized eigenfunctions are $\mathbf{f}_1 = 1$, $\mathbf{f}_2 = \sqrt{2} \cos \pi(\cdot)$ and $\mathbf{f}_3 = \sqrt{2} \sin \pi(\cdot)$ with eigenvalues $\lambda_1 = \frac{1}{2}$, $\lambda_2 = \lambda_3 = \frac{1}{4}$. We take parameters $T = 1$, $\gamma = 0.3$, $A = 0.5$, $B = 0.6$, $D = 2$, $\sigma = 0.5$, $Q = 0.3$, $\Gamma = 2$, $R = 1.5$, $Q_f = 0.8$, $\Gamma_f = -0.8$. The numerical solutions of $(\Pi, P^\perp, P^{\frac{1}{2}}, P^{\frac{1}{4}})$ to Riccati equations (7) and (82)–(83) are given in Figure 3a. We partition $[0, 1]$ uniformly into 200 subintervals, and take their left endpoints as the values of α and β . The initial values ξ_α are taken from Gaussian distribution $\mathcal{N}(2, 0.1)$. Then the state process \bar{x}_α in (34) with 200 different α values in the infinite population model and the corresponding control \bar{u}_α are shown in Figures 1b–1c. In addition, to fully capture the heterogeneity among agents across different nodes, we plot the graphon section g_α for various values of α in Figure 2a, and display the corresponding state and control trajectories in Figures 2b–2c. As shown in Figure 2a, the curves of graphon sections $g(0.25, \cdot)$ and $g(0.75, \cdot)$ are symmetric about the line $\alpha = 0.5$. Since the initial state means are the same, then the curves of $(z_{0.25}, S_{0.25})$ and $(z_{0.75}, S_{0.75})$ coincide in Figures 3b–3c.

7 Conclusion

This paper investigates a class of risk-sensitive GMFGs, for which decentralized strategies are determined from the GMFG equation system as a family of fully coupled forward-backward equations indexed by the nodal parameter. The unique solvability of the GMFG equation system is established using both the fixed point method and the method of continuity.

For performance analysis of the resulting decentralized strategies, novel exponential type error estimates are employed to establish an ε -Nash equilibrium theorem.

Our model does not include common noise in the agent dynamics since we are primarily interested in decentralized state feedback strategies while such strategies are generally inadequate for the search of an asymptotic Nash equilibrium when common noise appears.




Figure 1: Graphon limit, the trajectories of state and control for 200 different α values under the graphon limit.




Figure 2: Graphon section, the trajectories of the state and control with 4 different α values.




Figure 3: The solutions to (i) Riccati equations, and (ii) the GMFG equation system (18).

References

- [1] H. Abou-Kandil, G. Freiling, V. Ionescu, and G. Jank. Matrix Riccati equations in control and systems theory. Springer Basel AG, Basel, Switzerland, 2003.
- [2] A. Aurell, R. Carmona, and M. Lauriere. Stochastic graphon games: II. the linear-quadratic case. Appl. Math. Optim., 85(3):39, 2022.
- [3] T. Başar and G. J. Olsder. Dynamic noncooperative game theory, volume 23. SIAM, Philadelphia, PA, 1998.
- [4] D. Bauso, X. Zhang, and A. Papachristodoulou. Density flow in dynamical networks via mean-field games. IEEE Trans. Autom. Control, 62(3):1342–1355, 2017.
- [5] E. Bayraktar, S. Chakraborty, and R. Wu. Graphon mean field systems. Ann. Appl. Probab., 33(5):3587–3619, 2023.

- [6] A. Bensoussan, J. Frehse, and H. Nagai. Some results on risk-sensitive control with full observation. *Appl. Math. Optim.*, 37:1–41, 1998.
- [7] A. Bensoussan, J. Frehse, and P. Yam. Mean field games and mean field type control theory, volume 101. Springer, New York, USA, 2013.
- [8] C. Borgs, J. T. Chayes, L. Lovász, V. T. Sós, and K. Vesztegombi. Convergent sequences of dense graphs I: Subgraph frequencies, metric properties and testing. *Adv. Math.*, 219(6):1801–1851, 2008.
- [9] C. Borgs, J. T. Chayes, L. Lovász, V. T. Sós, and K. Vesztegombi. Convergent sequences of dense graphs II. Multiway cuts and statistical physics. *Ann. Math.*, pages 151–219, 2012.
- [10] P. E. Caines and M. Huang. Graphon mean field games and their equations. *SIAM J. Control Optim.*, 59(6):4373–4399, 2021.
- [11] P. Cardaliaguet, F. Delarue, J. M. Lasry, and P. L. Lions. The master equation and the convergence problem in mean field games:(AMS-201). Princeton University Press, 2019.
- [12] R. Carmona, D. B. Cooney, C. V. Graves, and M. Lauriere. Stochastic graphon games: I. the static case. *Math. Oper. Res.*, 47(1):750–778, 2022.
- [13] R. Carmona and F. Delarue. Probabilistic theory of mean field games with applications I–II. Springer, New York, USA, 2018.
- [14] Y. Chen, T. Li, and Z. Xin. Risk-sensitive mean field games with major and minor players. *ESAIM-Control Optim. Calc. Var.*, 29:6, 2023.
- [15] S. N. Cohen and R. J. Elliott. Stochastic calculus and applications, volume 2. Springer, New York, 2015.
- [16] F. Delarue. Mean field games: A toy model on an Erdős-Renyi graph. *ESAIM. Proc. Surv.*, 60:1–26, 2017.
- [17] T. E. Duncan. Linear-exponential-quadratic Gaussian control. *IEEE Trans. Autom. Control*, 58(11):2910–2911, 2013.
- [18] G. E. Espinosa and N. Touzi. Optimal investment under relative performance concerns. *Math. Financ.*, 25(2):221–257, 2015.
- [19] W. H. Fleming and D. Hernández-Hernández. Risk-sensitive control of finite state machines on an infinite horizon I. *SIAM J. Control Optim.*, 35(5):1790–1810, 1997.
- [20] W. H. Fleming and W. M. McEneaney. Risk-sensitive control on an infinite time horizon. *SIAM J. Control Optim.*, 33(6):1881–1915, 1995.
- [21] R. Foguen-Tchuendom, S. Gao, P. E. Caines, and M. Huang. Infinite horizon LQG graphon mean field games: Explicit Nash values and local minima. *Syst. Control Lett.*, 187:105780, 2024.
- [22] G. Fu and C. Zhou. Mean field portfolio games. *Financ. Stoch.*, 27(1):189–231, 2023.
- [23] S. Gao, P. E. Caines, and M. Huang. LQG graphon mean field games: Analysis via graphon-invariant subspaces. *IEEE Trans. Autom. Control*, 68(12):7482–7497, 2023.
- [24] R.A. Howard and J.E. Matheson. Risk-sensitive Markov decision processes. *Manag. Sci.*, 18(7):356–369, 1972.
- [25] Y. Hu and S. Peng. Solution of forward-backward stochastic differential equations. *Probab. Theory Related Fields*, 103:273–283, 1995.
- [26] M. Huang, P. E. Caines, and R. P. Malhamé. Large-population cost-coupled LQG problems with nonuniform agents: Individual-mass behavior and decentralized ϵ -Nash equilibria. *IEEE Trans. Autom. Control*, 52(9):1560–1571, 2007.
- [27] M. Huang, R.P. Malhamé, and P.E. Caines. Large population stochastic dynamic games: Closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle. *Commun. Inf. Syst.*, 6(1):221–252, 2006.
- [28] M. Huang and M. Zhou. Linear quadratic mean field games: Asymptotic solvability and relation to the fixed point approach. *IEEE Trans. Autom. Control*, 65(4):1397–1412, 2020.
- [29] D. Jacobson. Optimal stochastic linear systems with exponential performance criteria and their relation to deterministic differential games. *IEEE Trans. Autom. Control*, 18(2):124–131, 1973.
- [30] D. Lacker. A general characterization of the mean field limit for stochastic differential games. *Probab. Theory Relat. Field*, 165:581–648, 2016.
- [31] J. M. Lasry and P. L. Lions. Mean field games. *Jpn. J. Math.*, 2(1):229–260, 2007.
- [32] A. E. B. Lim and X. Y. Zhou. A new risk-sensitive maximum principle. *IEEE Trans. Autom. Control*, 50(7):958–966, 2005.

- [33] H. Liu, D. Firoozi, and M. Breton. LQG risk-sensitive single-agent and major-minor mean field game systems: A variational framework. *SIAM J. Control Optim.*, in press, 63(4):2251–2281, 2025.
- [34] L. Lovász. Large networks and graph limits, volume 60. American Mathematical Society, 2012.
- [35] L. Lovász and B. Szegedy. Limits of dense graph sequences. *J. Comb. Theory Ser. B*, 96(6):933–957, 2006.
- [36] A.M. Mathai and S.B. Provost. Quadratic forms in random variables: Theory and applications. Marcel Dekker, INC, New York, USA, 1992.
- [37] J. Moon and T. Başar. Risk-sensitive mean field games via the stochastic maximum principle. *Dyn. Games Appl.*, 9(4):1100–1125, 2019.
- [38] Z. Pan and T. Başar. Model simplification and optimal control of stochastic singularly perturbed systems under exponentiated quadratic cost. *SIAM J. Control Optim.*, 34(5):1734–1766, 1996.
- [39] A. Pazy. Semigroups of linear operators and applications to partial differential equations, volume 44. Springer Science & Business Media, New York, USA, 2012.
- [40] S. Peng and Z. Wu. Fully coupled forward-backward stochastic differential equations and applications to optimal control. *SIAM J. Control Optim.*, 37(3):825–843, 1999.
- [41] N. Saldi, T. Başar, and M. Raginsky. Approximate Markov-Nash equilibria for discrete-time risk-sensitive mean-field games. *Math. Oper. Res.*, 45(4):1596–1620, 2020.
- [42] R. E. Showalter. Monotone operators in Banach space and nonlinear partial differential equations, volume 49. American Mathematical Society, 2013.
- [43] H. Tembine, Q. Zhu, and T. Başar. Risk-sensitive mean-field games. *IEEE Trans. Autom. Control*, 59(4):835–850, 2014.
- [44] B. Wang and M. Huang. Mean field production output control with sticky prices: Nash and social solutions. *Automatica*, 100:90–98, 2019.
- [45] Y. Wang and M. Huang. Risk-sensitive linear-quadratic mean-field games: Asymptotic solvability and decentralized $O(1/N)$ -Nash equilibria. *J. Syst. Sci. Complex.*, 38(1):436–459, 2025.
- [46] P. Whittle. Risk-sensitive optimal control. American Mathematical Society, New York, USA, 1990.